

A TAXONOMIC REVISION OF THE GENUS
PROCOLOPHON AND THE
PHYLOGENETIC RELATIONSHIPS OF
PROCOLOPHONOID REPTILES

Juan Carlos Cisneros Martínez

A thesis submitted to the Faculty of Science, University of the Witwatersrand, in
fulfilment of the requirements for the degree of Doctor of Philosophy

Johannesburg, 2006


CONTENTS	Page
DECLARATION	ii
ABSTRACT	iii
ACKNOWLEDGEMENTS	iv
PAPERS	v
LIST OF FIGURES	vi
1 INTRODUCTION	1
1.1 Overview on procolophonids and related taxa	1
1.2 Distribution of procolophonids in Pangaea	6
1.3 Previous studies on <i>Procolophon</i>	9
2 MATERIALS AND METHODS	12
2.1 Specimens	12
2.2 Preparation	13
2.3 Illustration	13
2.4 Phylogenetic analyses	14
3 PAPER: A PROCOLOPHONOID REPTILE WITH TEMPORAL FENESTRATION FROM THE MIDDLE TRIASSIC OF BRAZIL	15
3.1 Introduction	16
3.2 Material and methods	17
3.3 Systematic palaeontology	17
3.4 Description	18
3.5 Discussion	22
3.6 References	25
3.7 Appendix A: Phylogenetic characters	29
3.8 Appendix B: Data matrix	30
4 PAPER: TAXONOMIC STATUS OF THE TRIASSIC REPTILE PROCOLOPHON IN GONDWANA	34
4.1 Introduction	35
4.2 Taxonomic history	36
4.3 Comparisons and discussion	38

4.3.1	The genus <i>Procolophon</i> in the Paraná Basin	38
4.3.2	Specimens with temporal fenestrae from the Karoo Basin	40
4.3.3	The genus <i>Procolophon</i> in the Transantarctic Mountains	44
4.4	Systematic palaeontology	46
4.5	Biogeographical considerations	49
4.6	Conclusions	49
4.7	References	51
4.8	Appendix	57
5	PAPER: PHYLOGENETIC RELATIONSHIPS OF	
	PROCOLOPHONID PARAREPTILES	68
5.1	Introduction	70
5.2	Taxonomic background and previous analyses	71
5.2.1	Nomenclatural remarks	73
5.3	Analysis and methods	74
5.3.1	Ingroup and outgroup	75
5.4	Character description	76
5.4.1	Cranium surface	77
5.4.2	Palate and braincase	82
5.4.3	Mandible	84
5.4.4	Dentition	84
5.4.5	Axial skeleton	88
5.4.6	Appendicular skeleton	89
5.4.7	Dermal ossifications	91
5.5	Results	92
5.5.1	Clade A: Procolophonidae Lydekker 1890	92
5.5.2	Clade E:	93
5.5.3	Clade F: Theledectinae new taxon	94
5.5.4	Clade H: “horned procolophonids”	95
5.5.5	Clade I: Procolophoninae Lydekker 1890	95
5.5.6	Clade K	96
5.5.7	Clade N: Leptopleuroninae Ivakhnenko 1979	97
5.5.8	Clade P	98

5.5.9	General remarks	98
5.6	Evolutionary history and feeding ecology	101
5.7	Biostratigraphic considerations	103
5.7.1	The Permian record of procolophonids	103
5.7.2	The Ladinian-Early Carnian record of procolophonids	107
5.8	Conclusions	108
5.9	References	110
5.10	Appendix: Data matrix	121
6	PAPER: A BASAL PROCOLOPHONID REPTILE FROM THE LOWER TRIASSIC OF SOUTH AFRICA	132
6.1	Introduction	133
6.2	Systematic palaeontology	134
6.3	Description	135
6.4	Discussion	140
6.5	References	142
7	DISCUSSION	151
7.1	Considerations on morphology and evolution	151
7.2	Biostratigraphic remarks	152
7.2.1	The appearance of bicuspid dentition	152
7.2.2	The age of the Sanga do Cabral Formation	154
7.2.3	The Ladinian-Carnian hiatus of procolophonids	154
7.3	Biogeography	155
8	CONCLUSIONS	157
9	REFERENCES	158
	APPENDIX: PUBLISHED PAPERS AND PARALLEL PRODUCTION	178

I declare that this thesis is my own work. It is being submitted for the Degree of Doctor of Philosophy in the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination in any other university.

I state that the papers contained in this thesis are my own, unaided work, except for the paper “A procolophonoid reptile with temporal fenestration from the Middle Triassic of Brazil”, written in collaboration. In this work, I contributed to the anatomical description and discussion, and undertook all the illustrations of the taxon.



24th of April 2007

ABSTRACT

This study presents a taxonomic revision of some procolophonoid parareptiles and a detailed, global analysis of procolophonid intrarelationships. The poorly known genus *Candelaria*, from the Middle Triassic of Brazil, is identified on the basis of new material as an owenettid, rather than a procolophonid as previously thought. Thus, *Candelaria* represents the youngest owenettid and the first member of this group from South America. The cranium of *Candelaria* is also remarkable for having temporal fenestrae, and the significance of this character within the Parareptilia is discussed. Based on a comprehensive review of specimens referred to different *Procolophon* species, it is proposed that only the type species, *Procolophon trigoniceps*, is valid. Thus, *Procolophon* specimens from Brazil, South Africa, and Antarctica are all referable to *P. trigoniceps*. Consequently, *P. trigoniceps* has one of the broadest known geographic ranges among Triassic tetrapod species.

A comprehensive cladistic analysis of procolophonids more firmly resolves the relationships within that group. The analysis reveals that Procolophoninae and Leptopleuroninae are valid monophyletic groups, whereas Spondylolestinae is paraphyletic. The species formerly assigned to the genus '*Thelegnathus*' from the Middle Triassic of South Africa, and those assigned to '*Eumetabolodon*' from the Lower-Middle Triassic of China, are paraphyletic.

The poorly known *Spondylolestes* from the *Dicynodon* Assemblage Zone of South Africa is considered valid and possibly represents the only Permian procolophonid in Gondwana. A new species, *Kitchingnathus untabeni*, is identified in the *Lystrosaurus* Assemblage Zone of South Africa. It is a basal member of the Procolophonidae and co-occurs with *Procolophon* in the Upper Katberg Formation. The new taxon is characterized by the presence of a large number of thin, bicuspid teeth. Character optimisation indicates that bicuspid teeth were acquired independently in *K. untabeni*, and hence originated twice during

procolophonid evolution. A review of procolophonid records worldwide reveals a fossil hiatus for members of this group in the Ladinian and most of the Carnian.

ACKNOWLEDGEMENTS

I wish to thank my supervisor Bruce S. Rubidge for his unconditional support and encouragement during three years of hard work at the Bernard Price Institute for Palaeontological Research in Johannesburg. A grant from the Palaeontological Scientific Trust in Johannesburg made possible my dedication to this project. I am grateful to my co-supervisor Ross Damiani for his advice during this study and to Rainier Schoch for providing facilities at the Staatliches Museum für Naturkunde, Stuttgart, where the final stages of this work were completed. Both are acknowledged for their hospitality in Stuttgart.

The following people are acknowledged for fossil access during this study: Alex Cook and Kristen Spring (Queensland Museum), Átila da Rosa (Universidade Federal de Santa Maria), Billy de Klerk (Albany Museum), Cesar Schultz (Universidade Federal do Rio Grande do Sul), Elize Butler and Jennifer Botha (National Museum), Fang Zheng and Li Jinling (Institute of Vertebrate Palaeontology and Palaeoanthropology), Mike Raath (Bernard Price Institute for Palaeontological Research), Roger Smith and Sheena Kaal (Iziko South African Museum) and the Rubidge family (Rubidge Collection). Doctor M. Mbense, Charlton Dube (Bernard Price Institute for Palaeontological Research) and John Nyaphuli (National Museum) are acknowledged for their outstanding preparation.

Special thanks to my colleagues Sally Reynolds, Fernando Abdala and Adam Yates for their support, advice and happy moments during my stay in South Africa. This work would have not been possible without tons of dark chocolate and inspiration from the music of Tangerine Dream, Klaus Schulze, Laurie Anderson, Kraftwerk, Jean Michel Jarre, Dead Can Dance, Erik Satie, Wolfgang Amadeus Mozart and Johann Sebastian Bach.

LIST OF PAPERS SUBMITTED AS PART OF THIS THESIS

This thesis is presented in the format of a compilation of four papers which were published/submitted by the author while registered for a PhD degree.

Cisneros, J.C., Damiani, R., Schultz, C. Rosa, A. da, Schwanke, C., Neto, L.W. and Aurélio, P.L.P. (2004) A procolophonoid reptile with temporal fenestration from the Middle Triassic of Brazil. *Proceedings of the Royal Society of London, Series B, Biological Sciences*, vol. 271, pp. 1541-1546.

Cisneros, J.C. Taxonomic status of the Triassic reptile *Procolophon* in Gondwana. *Palaeontologia africana* (in press)

Cisneros, J.C. Phylogenetic relationships of procolophonid parareptiles. *Journal of Systematic Palaeontology* (submitted)

Cisneros, J.C. A basal procolophonid reptile from the Lower Triassic of South Africa. *Palaeontology* (submitted)

LIST OF FIGURES

Chapter 3

Figure 1. Reconstruction of the skull of the owenettid parareptile *Candelaria barbouri* (Price, 1947) from the Middle Triassic of Brazil, based on UFSM 11076, UFSM 11131 and DGM 314R. (a) Dorsal view; and (b) left lateral view.

Characters 1-5 relate to the temporal fenestra and are listed in the diagnosis.

Abbreviations: a, angular; ar, articular; c, coronoid process; d, dentary; f, frontal, j, jugal; la, lacrimal; m, maxilla; n, nasal; p, parietal; pf, prefrontal; pm, premaxilla; po, postorbital; pof, postfrontal; q, quadrate; qj, quadratojugal; sa, surangular; sq, squamosal; st, supratemporal..... 31

Figure 2. Stratocladogram of procolophonoid interrelationships, including ghost lineages (white extensions of black bars). The Owenettidae is a monophyletic taxon that includes *Candelaria*, while the Early Triassic taxa *Coletta* and *Sauropareion* are transitional forms between the Owenettidae and Procolophonidae. The phylogeny is based on a PAUP 3.1.1 (Swofford, 1993) analysis of the data matrix in Appendix 2, and represents one of two most parsimonious trees, chosen on the basis of greatest stratigraphic congruence, found using the branch-and-bound algorithm. Tree length=31 steps, consistency index=0.81, and rescaled consistency index=0.80. Biostratigraphy of South Africa from Rubidge (1995); southern Brazilian Triassic from Abdala *et al.* (2001); southern Brazilian Permian adapted from Langer (2000) and Malabarba *et al.* (2003). Hatchure indicates a sedimentary hiatus. Abbreviations: *Cist.*, *Cistecephalus*; *Cynog.*, *Cynognathus*; *Dicyn.*, *Dicynodon*; *Dinod.*, *Dinodontosaurus*; 'Endo.', 'Endothiodon'; *Lyst.*, *Lystrosaurus*; *Procol.*, *Procolophon*; *Travers.*, *Traversodontid*; *Trop.*, *Tropidostoma*..... 32

Figure 3. Simplified cladogram of 'anapsid' reptiles, showing distribution (in bold face) of all known taxa which possess temporal fenestrae (see discussion). Skull outlines, from left to right, are: *Stereosternum tumidum*, *Millerosaurus nuffieldi*,

Macroleter poezicus, *Tokosaurus perforatus*, *Bradysaurus baini*, *Procolophon laticeps*, *Sauropareion anoplus*, *Owenetta kitchingorum*, *Candelaria barbouri*, *Lanthanosuchus watsoni* and *Acleistorhinus pteroticus*; temporal fenestrae are shaded in black. Phylogeny adapted from Reisz and Scott (2002). Drawings not to scale..... 33

Chapter 4

Figure 1. Life reconstruction of *Procolophon trigoniceps*. Note the presence of cheeks; quadratojugal processes covered by long keratinous spines; and large digging-claws. The skeleton of *Procolophon* entirely supports these features (Carroll and Lindsay, 1985; deBraga, 2003). Numerous minor keratinous spines are common over large or stocky-bodied modern lizards; these structures do not leave traces in the skeleton (e.g. *Iguana*, *Phrynosoma*, *Uromastix*; pers. obs.). The long quadratojugal and supratemporal spines of *Procolophon* may have acted as an anti-predatory mechanism, as in phrynosomatid lizards (Young *et al.*, 2004)..... 58

Figure 2. BMNH R1726 holotype of *Procolophon trigoniceps*, cranium in dorsal and left views..... 59

Figure 3. A, MCN PV1904 holotype of *Procolophon brasiliensis*, cranium in palatal view. B, UFRGS PV231T holotype of *Procolophon pricei*, cranium in palatal view. Scale bar represents 10 mm..... 60

Figure 4. Palate of *Procolophon trigoniceps* specimens from the South African Karoo. A, CGP 1-89; B, BP/1/4014; C, BP/1/4248; D, NM QR1447; E, AMNH 5693; F, BP/1/966. Arrows in C and F indicate posterior enlarged vomerine teeth. Arrows in D and E indicate the last tooth in the pterygoid-palatine tooth row. The mandible is in occlusion in BP/1/4248, NM QR1447 and AMNH 5693. Scale bar is 10 mm for A-E and 17 mm for F..... 61

Figure 5. *Procolophon* specimens with temporal fenestrae. A, CGP 1-127; B, BMNH R1949; C, BMNH R 3583 holotype of *P. laticeps*. Scale bar represents 10 mm, arrows indicate temporal fenestrae..... 62

Figure 6. Temporal fenestrae in *Procolophon*. A-D, CGP 1-127. E, F, BMNH R3583 holotype of *P. laticeps*. G-J, BMNH R1949. A, C, E, G, I, left lateral views. B, D, F, H, J, right lateral views..... 62

Figure 7. Paroccipital process of the right opisthotic in *Procolophon*, in lateral view. A. BMNH R1949. B, CGP 1-127. C, BMNH R4087. A and B are individuals with temporal openings. Scale bar represents 5 mm..... 64

Figure 8. AMNH 9506, *Procolophon trigoniceps* from Shackleton Glacier, Transantarctic Mountains. A, skeleton in dorsal view. B, detail of palate in ventral view, arrow points to the depression between the vomers. Scale bars represent 5 mm (A) and 2 mm (B)..... 65

Figure 9. Comparison between A, *Teratophon spinigenis* (BP/1/4587), B, *Eumetabolodon bathycephalus* (IVPP V6064) and C, *Procolophon trigoniceps* (BP/1/5927b); showing differences in dentition. Scale bar represent 5 mm for A, B and 3.5 mm for C..... 66

Figure 10. *Procolophon trigoniceps*, silicone impression of AM 358, a left hind limb in flexor aspect. The arrow in black shows the position where the fifth metatarsal should be located. The excellent preservation of this limb indicates that the absence of the fifth metatarsal in *Procolophon* is a natural feature rather than a post-mortem phenomenon as suggested by deBraga (2003). Although this specimen is smaller than other *Procolophon* individuals, all bones are well ossified. Scale bar is 5 mm..... 67

Chapter 5

Figure 1. Crania of procolophonids in dorsal (A-E) and right lateral (F-J) view, showing some characters used in this study (states indicated in brackets). A, *Tichvinskia vjatkensis* (PIN 954/1); B, *Procolophon trigoniceps* (NM QR3201); C, *Neoprocolophon asiaticus* (IVPP V866); D, *Pentaedrusaurus ordosianus* (IVPP V8735); E, *Hypsognathus fenneri* (YPM 55831); F, *Tichvinskia vjatkensis* (PIN 954/1); G, *Procolophon trigoniceps* (BMNH R4087); H, *Neoprocolophon asiaticus* (IVPP V866); I, *Pentaedrusaurus ordosianus* (IVPP V8735); J, *Hypsognathus fenneri* (YPM 55831). All drawn from originals except *Tichvinskia*, redrawn from Ivakhnenko (1979), presumably reconstructed. Note that the sutures in *Neoprocolophon* differ from Young (1957). Coding of character 5: A vertical line (a) is traced from the cranial roof to the alveolar margin, tangential to the anterior border of the orbitotemporal fenestra, and a line (b) is traced from (a) to the tip of the snout, perpendicular to (a). The snout is considered “long and flat” if (a) is shorter than (b); and “deep and short” if (a) is equal or longer than (b). Because the tip of the snout is not fully preserved in *Neoprocolophon*, coding of this character was based on a conservative estimation of the total length of the snout. Drawings not to scale..... 124

Figure 2. Maxillary teeth of procolophonids showing some characters from this study. A, *Coletta seca* (CGP/1/1003), left vii-ix in labial view, and right vii-ix in occlusal view (note that cusps are missing); B, *Arcadia* procolophonid (QMF 49510), left ii-iv in labial view, and left ii-iii in occlusal view; C, *Soturnia caliodon* (MCN PV2738) right ii-iii molariforms in lingual and occlusal views; D, *Procolophon trigoniceps* (SAM PK-K9998), right v-vii in labial (slightly posterior) and occlusal views. Drawings not to scale..... 126

Figure 3. Postcranial elements of procolophonoids, showing some characters from this study. A-C, manus; D-F, humerus, G-I, interclavicle, J-L, femur. *Barasaurus besairiei*: A, manus (SAM PK-K8275); D, right humerus, dorsal view (SAM PK-K-8282); G, interclavicle, ventral view (SAM PK-K8276); J, right femur, anterior

view (SAM PK-K-8282). *Procolophon trigoniceps* (BP/1/962): B, right manus, dorsal view; E, right humerus, ventral view; H, interclavicle, ventral view; K, right femur, anterior view. *Pentaedrusaurus ordosianus* (IVPP V8735): C, left manus, dorsal view; F, left humerus, ventral view (note that it differs from Li, 1989); I, interclavicle, ventral view; L, right femur, anterior view. Drawings not to scale..... 127

Figure 4. Single most parsimonious tree. Tree length=134 steps, consistency index (excluding uninformative characters) = 0.642, retention index = 0.796. DI and SR values are given next to each node (DI/SR, respectively). DI values were calculated from 6967 trees. SR was performed with 5000 replicates and 10 repetitions (p=0.33) under Traditional Search option (random addition sequences plus tree bisection-reconnection)..... 128

Figure 5. Single most parsimonious tree showing non-ambiguous synapomorphies for clades, and autapomorphies for terminal taxa. Reversals are indicated by white boxes..... 129

Figure 6. Single most parsimonious tree showing important feeding-related acquisitions in Procolophonids. A, thin conical tooth with circular base, medial (plesiomorphic state); B, tooth with labiolingually expanded base, mesial view; C, bulbous tooth, mesial view; D, bicuspid tooth, mesial view; E, multiple tooth rows; F, teeth inset from the lateral maxillary surface; G, molariforms with large occlusal areas; H, low articular; I, deep dentary; J, sharp, multiple cusped molariform; K, molariform with large occlusal area and a deep occlusal depression..... 130

Figure 7. Single most parsimonious tree and the geological record of procolophonoids. Black bars represent the stratigraphic ranges, dashed bars represent uncertain stratigraphic ranges. The global hiatus of procolophonid records in Ladinian-Lower Carnian rocks is represented in gray. The minimum divergence time (MDT) of the Procolophonidae from the Owenettidae predicts the

existence of a procolophonid as early as the Late Wuchiapingian. Stratigraphic ranges of taxa and geological dates based on: Li (1989), Lucas (1999), Smith (2000), Spencer and Benton (2000), Sues *et al.* (2000), Modesto *et al.* (2002), Cisneros *et al.* (2004), Rubert and Schultz (2004), Golubev (2005), Sennikov and Golubev (2005) and Hancox *et al.* (in press). SGCS (Standard Global Chronostratigraphic Scale) stages after Gradstein and Ogg (2004)..... 131

Figure 8. Cranium of *Spondylolestes rubidgei*, RC 3, holotype. **A**, dorsal; **B**, ventral; **C**, right lateral; and **D**, left lateral views of the skull. **E** and **F**, fragments of left maxilla and dentary in occlusion; **E**, dorsal view, showing maxillary dentition in basal cross-section; and **F**, lingual view showing both dentitions. Abbreviations: **ap**, ascendent process of the premaxilla; **d**, dentary; **f**, frontal; **iv**, interpterigoyd vacuity; **mx**, maxilla. Arrows indicate correspondence among maxillary teeth. Scale bar represents 5 mm for **A-D**, and 10 mm for **E** and **F**... 132

Chapter 6

Figure 1. **A**, *Kitchingnathus untabeni* n. g. n. sp., Lower Triassic, South Africa, BP/1/1187, holotype, photograph of the skeleton; **B**, outline of the skeleton. Abbreviations: ac, anterior coracoid; c, clavicle; cr, cranium; g, gastralia; i, interclavicle; il, ilium; f, femur; lm, left mandibular ramus; lmx, left maxilla; mp, metapodial; of, orbitotemporal fenestra; r, rib; rm, right mandibular ramus; sc, sacrum; vt, vertebra. Digits are identified with Roman numerals ii-v. Postcranial bones that could not be identified are not labelled..... 147

Figure 2. *Kitchingnathus untabeni* n. g. n. sp., Lower Triassic, South Africa, BP/1/1187, holotype. **A**, cranium, right view. **B**, right mandible in lateral view and selected teeth in occlusal view. **C**, left mandible in medial view and selected teeth in occlusal view. Abbreviations: amf, anterior maxillary foramen; d, dentary; cr, coronoid; f, frontal; j, jugal; l, lacrimal; mx, maxilla; n, nasal; op, opisthotic; p, parietal; pf, pineal foramen; pmx, premaxilla; pof, postfrontal; prf, prefrontal; pt, pterygoid; q, quadrate; qj, quadratojugal; sp, splenial; st, supratemporal; ect,

ectopterygoid. Unshaded surfaces represent features preserved as natural casts. Roman numerals indicate tooth positions. Scale bar represents 5 mm, except for dentitions in occlusal view where it represents 2 mm..... 148

Figure 3. *Kitchingnathus untabeni* n. g. n. sp., BP/1/1187. Photographs of lower dentition. A, right mandible, lateral view. B, left mandible, dorsolateral view. C, left mandible, posterodorsal view..... 149

Figure 4. *Kitchingnathus untabeni* n. g. n. sp., BP/1/1187. Reconstruction of the skull in lateral view. Abbreviations: an, angular; ar, articular; bo, basioccipital; d, dentary; cr, coronoid; ect, ectopterygoid; f, frontal; j, jugal; l, lacrimal; mx, maxilla; n, nasal; op, opisthotic; p, parietal; pf, pineal foramen; pmx, premaxilla; po, postorbital; pof, postfrontal; prf, prefrontal; pt, pterygoid; q, quadrate; qj, quadratojugal; sa, surangular; sq, squamosal; st, supratemporal; ect, ectopterygoid..... 150

Figure 5. Simplified cladogram of procolophonid relationships. The presence of bicuspid marginal teeth is indicated by closed circles. For details see Chapter 5..... 151

Chapter 7

Figure 1. Composite cladogram of the Procolophonoidea, adapted from the phylogenetic analyses performed in chapters 3 and 6. *Spondylolestes rubidgei* and *Kinelia broomi* were not included in the cladistic analyses and their phylogenetic positions are tentative..... 157