





Supporting maternal and child nutrition: views from community members in rural Northern Ghana

Cornelius Debpuur¹, Engelbert A Nonterah^{1,2} , Samuel T Chatio^{1,*}, James K Adoctor¹, Edith Dambayi¹, Paula Beeri¹, Esmond W Nonterah¹, Doreen Ayi-Bisah¹, Daniella Watson³ , Sarah H Kehoe⁴, Maxwell A Dalaba¹, Winfred Ofofu⁵, Raymond Aborigo¹, Paul Welaga¹, Abraham R Oduro¹, Marie-Louise Newell^{3,6} and Mary Barker^{4,7} for the INPreP Study Group

¹Clinical Science Department, Navrongo Health Research Centre, Ghana Health Service, Hospital Road, 00233 Navrongo, Ghana: ²Julius Global Health, Julius Centre for Health Science and Primary Care, University Medical Centre Utrecht, Utrecht University, Utrecht, the Netherlands: ³Department of Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK: ⁴Medical Research Council Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK: ⁵Upper East Regional Health Directorate, Ghana Health Service, Bolgatanga, Ghana: ⁶Faculty of Health Sciences, School of Public Health, University of the Witwatersrand, Johannesburg, South Africa: ⁷National Institute of Health Research, Southampton Biomedical Research Centre, University Hospital Southampton National Health Service Foundation Trust, Southampton, UK

Submitted 25 December 2019: Final revision received 2 July 2020: Accepted 24 July 2020: First published online 25 September 2020

Abstract

Objective: Despite efforts to improve maternal and child nutrition, undernutrition remains a major public health challenge in Ghana. The current study explored community perceptions of undernutrition and context-specific interventions that could improve maternal and child nutrition in rural Northern Ghana.

Design: This exploratory qualitative study used ten focus group discussions to gather primary data. The discussions were recorded, transcribed and coded into themes using Nvivo 12 software to aid thematic analysis.

Setting: The study was conducted in rural Kassena-Nankana Districts of Northern Ghana.

Study participants: Thirty-three men and fifty-one women aged 18–50 years were randomly selected from the community.

Results: Most participants reported poverty, lack of irrigated agricultural land and poor harvests as the main barriers to optimal nutrition. To improve maternal and child nutrition, study participants suggested that the construction of dams at the community level would facilitate all year round farming including rearing of animals. Participants perceived that the provision of agricultural materials such as high yield seedlings, pesticides and fertiliser would help boost agricultural productivity. They also recommended community-based nutrition education by trained health volunteers, focused on types of locally produced foods and appropriate ways to prepare them to help improve maternal and child nutrition.

Conclusion: Drawing on these findings and existing literature, we argue that supporting community initiated nutrition interventions such as improved irrigation for dry season farming, provision of agricultural inputs and community education could improve maternal and child nutrition.

Keywords

Community interventions
Maternal and child nutrition
Community perceptions
Ghana

In Ghana, maternal and child undernutrition remains a major public health challenge and Ghana has been classified among the thirty-six countries global accounting for 90% of all stunting among children under 5 years^(1,2). According to the 2014 Ghana Demographic and Health Survey, about 20% of children under 5 years were

stunted⁽²⁾. In 2018, the prevalence rates of stunting were 37, 36, and 25% for the Northern, Upper East and Upper West regions, respectively, while the prevalence rates of wasting were 11, 11, and 7% in the same regions⁽³⁾. Among children under 5 years, 66% were anaemic, while 44% of women 15–49 years of age were anaemic⁽²⁾. Additionally,

*Corresponding author: Email schatio@yahoo.co.uk

© The Author(s), 2020. Published by Cambridge University Press on behalf of The Nutrition Society



41% of women nationally are now recorded as overweight and obese, suggesting a growing double burden of malnutrition. Urban Ghanaian women are increasingly becoming overweight and obese, while women in rural areas remain underweight⁽⁴⁾. In 2010, about 1.2 million people in the general population in Ghana were food insecure with a greater proportion of this group coming from the Northern part of the country⁽⁵⁾. Low productivity in the agriculture sector as a result of low soil fertility and unreliable rainfall has been a major problem for food and nutrition security in Ghana⁽⁵⁾.

Ghana's National Nutrition Policy aims to increase the coverage of high-impact nutrition-sensitive and nutrition-specific interventions to ensure optimal nutrition of Ghanaians, with special reference to improving maternal and child nutrition⁽⁶⁾. The policy is intended to educate people about the importance of investing in nutrition, guide the implementation of evidence-based nutrition interventions and promote healthy lifestyles and appropriate dietary habits⁽⁶⁾. Various interventions, such as postharvest food storage techniques and production of animal-sourced foods, have been introduced to increase and improve access to nutritious foods in sub-Saharan African countries including Ghana^(7,8). Integrated agriculture–nutrition interventions in Ghana have resulted in significantly higher child length and height *z* scores, improved diet diversity and quality of infant feeding practices, some of these linked to women growing empowerment^(9,10). Also, growth promotion, micronutrient supplementation, behaviour change communication about infant and young child feeding practices and community management of severe acute malnutrition have also been introduced to improve nutritional status⁽¹¹⁾.

However, access to nutritious foods remains a challenge in Ghana where factors including diseases and pests affect crop yields and flooding and poverty affect availability of and access to nutritious foods⁽¹²⁾. Climate change exacerbates problems of access to food in Ghana. It disrupts agricultural outputs, especially in the Northern region of Ghana which is particularly vulnerable to climate change, and where millions of poor smallholder farmers rely on rainfall for food and income for their families^(13,14).

Despite efforts, maternal and child undernutrition still remains a challenge in Ghana^(3,15). There is limited evidence to guide appropriate community-level initiatives to address nutritional challenges in developing countries such as Ghana. Designing interventions to improve nutrition requires a clear understanding of the personal and contextual factors affecting patterns of food choice and consumption, including social and psychological factors^(16,17). The current study explored community perceptions on contextual strategies to improve maternal and child nutrition in rural Kassena-Nankana Districts of Northern Ghana.

This is a sub-study of a larger National Institute for Health Research-funded international collaboration Improved

Nutrition Preconception Pregnancy Post-Delivery of partners based in Ghana, Burkina Faso, South Africa and the UK. Improved Nutrition Preconception Pregnancy Post-Delivery aims to develop supportive delivery of nutrition interventions in the 'first 1000 days plus', meaning preconception, pregnancy and the first 2 years of life. The community's perceptions of such interventions are presented in this special series^(18–20). These were collected to inform the design of interventions to improve maternal and child health and nutrition in these populations.

Methods

Study design

This qualitative study draws on data from ten focus group discussions (FGD) with community-dwelling adult men and women aged 18–50 years. The discussions were conducted between January and April 2019. Qualitative research is descriptive of the process and the meanings gained through words⁽²¹⁾, which helps in capturing the feelings, experiences and perceptions of individuals on the issue under investigation. The approach was deemed appropriate because our study aimed to gain a deeper understanding of local people perceptions of contextual solutions to improve maternal and child nutrition.

Study site

The study was conducted in the Kassena-Nankana East and West Districts of Northern Ghana by the Navrongo Health Research Centre. The Navrongo Health Research Centre operates the Navrongo Health and Demographic Surveillance System in the two districts. The districts cover an area of 1675 km² of Sahelian savannah with a population of about 153 000⁽²²⁾. The main languages spoken in the area are Kasem and Nankani. The population is predominantly rural with subsistence farming as the mainstay of the economy. There are two distinct seasons: the rainy season from May to October and the dry season from November to April. The population largely lives on subsistence crops including millet, sorghum, rice, maize and groundnuts⁽²³⁾. Fruits and vegetables such as tomatoes, onions, pepper, sweet potatoes, cabbage and lettuce are also produced in the area. Inhabitants of the two districts mostly live in multi-household compounds.

Sampling techniques

Data from the Navrongo Health and Demographic Surveillance System were used as the sampling frame. For data collection purposes, the Navrongo Health and Demographic Surveillance System area has been divided into five zones (East, West, North, South and Central)⁽²⁴⁾. The East and South zones are predominantly Nankani speaking while the West, Central and North are Kasem speaking zones. These zones are further divided into clusters. Two zones, one each in the Kasem (North) and Nankani (South)



speaking areas, were randomly selected for the study. Ten clusters (five in each zone) were randomly selected as communities where the FGD were conducted. In each cluster, about twenty individuals who met the age and sex criteria were randomly selected and were contacted by the data collectors. The first twelve people who gave consent were invited to participate in each FGD.

Training and data collection procedures

Four university graduates (research officers) with experience in conducting qualitative interviews were recruited and trained for data collection. A pre-test was conducted at the end of the training to evaluate the performance of data collectors and help finalise the discussion guides for data collection.

Each FGD was conducted by two research officers, one serving as a moderator and the other as an observer and a note taker. The data collectors visited individuals at home and invited them to participate in the FGD. The FGD were constituted separately based on gender (men, women), ethno-linguistics group (Kasem and Nankani) and further disaggregated by age (18–25, 26–39 and 40–50 years for women; 24–34 and 35–50 years for men). This enabled us to solicit views across different age–gender groups among the two main ethnic groups in the study area. A total of ten FGD (four with men and six with women) were conducted. A suitable venue was selected by study participants at the community level where the discussions were held using the two main local languages (Kasem and Nankani) and tape recorded with the consent of study participants. On average, the FGD lasted 1 h. A harmonised discussion guide was developed for all three study sites with enough flexibility to allow researchers to tailor the discussion to the local context. The guide covered areas such as maternal and child health issues, maternal and child nutritional problems and suggestions to improve maternal and child nutrition.

Data analysis techniques

The recordings were transcribed verbatim into English by graduate research officers who were native speakers with experience in transcribing qualitative interviews. A codebook was developed using the original research questions and themes that emerged from the data to guide data coding and analysis. The data were organised using QSR Nvivo 12 software for thematic analysis⁽²⁵⁾. The transcripts were initially coded independently by two members of the research team. The coding process involved a critical review of each transcript to identify emerging themes from the data and also based on the objectives of the study. The two coders then met to compare their independently identified themes. They resolved any divergence by re-reading the relevant sections of the transcripts together and agreed on the best fit interpretation of the data. The findings are presented in themes and supported by relevant quotes from the data.

Results

Socio-demographic characteristics of participants

Eighty-three adults took part in the study, of whom 49.7% were between 26 and 39 years. The majority of the participants were married and 51.8% were from the Kasem ethnic group. About 51.0% had between primary and junior high education with only 22.9% having secondary or higher qualification. About 45.0% of the participants were farmers. Almost 51.0% were unable to indicate their household monthly income because they did not know how much they earned (Table 1).

Community perspectives of factors affecting maternal and child nutrition

Study participants highlighted four main factors affecting maternal and child nutrition in the area. These comprised poverty, poor harvest, lack of irrigated agricultural land and perceived lack of support from men.

Poverty

Poverty was perceived to be a significant determinant of participants' nutrition across all ages and genders. Most participants believed poverty to be the main economic determinant limiting their access to nutritious food. They attributed this to the high rates of unemployment in the area. Study participants associated the paucity of job opportunities with lack of money to buy the food that would enhance their nutritional status.

Table 1 Socio-demographic characteristics of participants

Variable	Frequency	Percentage
Age (years)		
18–25	20	24.8
26–39	42	49.7
40+	21	25.5
Sex		
Male	32	38.6
Female	51	61.4
Ethnicity		
Kasem	43	51.8
Nankam	40	48.2
Marital status		
Never married	13	15.7
Married	66	79.5
Widowed	4	4.8
Educational status		
No education	22	26.5
Primary/junior high	42	50.6
Senior high/tertiary	17	22.9
Occupation		
Unemployed	13	15.7
Civil/public servant	3	3.6
Self employed	30	36.1
Farmer	37	44.6
Household monthly income (\$)		
19–72	29	35.0
74.3–129	7	8.4
130+	5	6.0
Unable to tell	42	50.6



The hand is less (meaning there is no money). To be honest, in this our community, money is a problem for us. There is no work for us to do to get money and support ourselves. It is only sufferings and hunger because when we need something we cannot buy because of poverty. (FGD-women-Kasem-18–25 years)

We know nutritious food helps the woman . . . but it is poverty that makes you unable to take care of the woman or buy nutritious food she needs to eat to help her and the children . . . (FGD-men-Nankani-35–50 years)

Poor harvest and lack of irrigated agriculture

Another challenge to good nutrition identified by study participants was poor harvest due to low soil fertility and unreliable rainfall. Both men and women in the discussions held the view that food production in the district was dictated by weather and often limited to the rainy season, which is short and unreliable. Study participants noted that farm harvests were affected by low soil fertility and diseases and pests that affect crop yields resulting in poor nutrition at the community level.

Our land is not good (infertile). We all farm but we don't get anything because there is no fertilizer. When it is raining season, we farm everywhere and we don't get returns on the farming because we don't have money to buy the medicine (pesticides) to spray and kill the diseases affecting our crops. (FGD-men-Kasem-35–50 years)

As for farming, there is nothing in there. You will work and work but you would not get anything there, the little you get from the farming you eat that. Within a short time, the food is finished . . . where will you get food to eat and be healthy? It is only sufferings. (FGD-women-Nankani-18–25 years)

The majority of the study participants were of the view that unavailability of water in the dry season made it difficult to grow vegetables and other food crops, thus restricting access to nutritious food in the community. They explained that vegetables were easily available during the rainy season, making it easier to enrich their diet, but this was not the case in the dry season.

. . . When it is raining season, we do gardening, but this dry season, there is no water for us to do the gardens or grow crops to get food during this time of the year. So as we are sitting, it is difficult for us to get nutritious food this time of the year. (FGD-men-Kasem-24–34 years)

Perceived lack of male support

Study participants also shared their views on how gender roles affect maternal and child nutrition in the area. Most of the women perceived that men were not taking up their

responsibilities of providing for the family. They recounted the difficulties they faced in the upbringing of their children, noting that the children's education, health needs and feeding were left under their care without support from their husbands. Study participants of both genders identified the lack of support by men as a significant barrier to access to nutritious foods by women and children as shown in the excerpts below:

To me, it is the drinking. The men, they married us and we are now suffering. They have money to buy alcohol and they know what to do outside and they have left the women. The problem is that they have refused to care for the children and now we (women) don't have anything to do and get money. How can we get nutritious food for the children? That is the problem we have in this community. (FGD-women-Nankani-26–39 years)

That is true, most men as soon as they get the money, they forget that they have women that they have to take care of let alone say children. Where will they (women and children) go and get food to eat if you the man is not concerned? So that is the problem for some of the men here. (FGD-men-Nankani-24–34 years)

Although the women did not directly suggest ways in which men could be more involved in family life, the men reflected on how they could support women, suggesting that they should be more patient and contribute to childcare.

Having patience, allow your temper to go down . . . you should know how to talk kindly to a pregnant woman. (FGD-men-Kasem-35–50 years)

The men too should have patience to bath the children, when they are sick. Give them their medications before they go out. (FGD-men-Kasem-35–50 years)

Proposed contextual interventions to improve community nutrition

Study participants identified various strategies that could improve maternal and child nutrition in the area. These included irrigated agricultural land, education on nutrition, provision of agricultural inputs and food supplements.

Irrigated agricultural land

A common suggestion among study participants was for a construction of dams in the communities to provide water for all year round agricultural activities. This would help community members to rear animals during the dry season and also make small gardens where they could grow nutritious foods and fresh vegetables to facilitate access to nutritious food for their families.

What I have to say is that if we are provided with water like dams for us to do gardens, we can get fresh



vegetables to eat during the dry season. This will be beneficial to us because it will help us get food. (FGD-woemn-Nankani-18–25 years)

... So, if we had water, we can get all the vegetables we need to eat and be healthy. With the water, we can grow trees, vegetables and when that happens, food will not be a problem for us in the community. I think that water can help us to do dry season gardens to enable us take care of our women and children. (FGD-men-Kasem-35–50 years)

Provision of agricultural materials

Some of the participants associated their poor farm yields with lack of agricultural materials. According to them, lack of these farming materials leads to poor yield. Participants were of the view that any support in terms of high yield seeds (groundnuts and beans) and also fertiliser and pesticides to spray insects that destroy their crops would improve their farm yields and thereby provide enough nutritious food for the community.

Most of us don't have ... So if we could get support in terms of some of the seedlings it will help us to have good harvest. (FGD-women-Kasem-40–50 years)

... Here, when you plant your crops and before you realize insects will eat everything. So I think if they are able to come and support us with chemicals (pesticides) and fertilizer at a reduced price that will enable us get good yield and there will be food for us to eat. (FGD-men-Nankani-35–50 years)

Some participants also wished for financial support to rear animals such as goats, sheep and fowls. They believed that meat from such animals could enrich the nutrition of their households. Moreover, they could sell some of the animals to buy other nutritious food that they did not produce to enhance their nutritional status.

What I want to be done here to help us especially the women is that if they could get support especially, financial support to rear goats and fowls, it will benefit us a lot. (FGD-women-Nankani-40–50 years)

If the youth could be supported to rear animals that would help the women in terms of their diet ... we could extract milk from cattle for the women to be drinking and for feeding the children. So if you could support us in rearing animals, it would help us. (FGD-men-Kasem-24–34 years)

Effective community education on nutrition

Study participants recommended that intensive education be provided to community members, especially women, on the types of locally produced nutritious foods they could eat and more importantly how to prepare them. To facilitate this approach, participants suggested that trained

community-based health volunteers could educate people in the communities on nutrition issues. This strategy was identified mainly by the men in the discussions as a way of improving community members' knowledge on nutrition and thereby contributing to enhance maternal and child nutritional status.

You see, now we don't have people to talk to us especially women about the types of foods especially the locally produced foods that is nutritious or what they are supposed to eat and be healthy. If they don't know, it would be difficult for them to eat good food or even how to prepare good food for their children. So if we could get people who will come and teach women, it would bring about massive change in this community. (FGD-men-Nankani-24–34 years)

... When you (refers to the research team) come and teach our people that will help. Also we have health volunteers in this community. We will select them to help this idea to work. That is something we have been yearning for, and we are not getting, so if we get such a thing we will support you because we know it is going to benefit us. (FGD-men-Kasem-35–50 years)

Food aid and nutrition supplementation

Some of the participants recalled several nutrition-specific interventions they had been offered in the past and felt that reintroducing these as well as improving their ability to farm would be good for the nutritional status of their women and children. These nutrition-specific interventions included providing food supplements during antenatal visits and at child welfare clinics.

... So for me if the health workers could give food items to pregnant women and say when you wake up prepare this and eat, take this for the children, use this to prepare food for the children. I think this can help. (FDG-men-Kasem-24–34 years)

At first they used to give us flour from the health facilities to pregnant women or mothers. They would give you the flour to go and prepare it for the child to eat and be strong and healthy. That is no more there at the health facilities, for a very long time. We don't remember when that was stopped. So when they do that again, it will be good. (FGD-women-Nankani-26–39 years)

Discussion

The current study explored community members' perceptions of factors affecting the nutrition of women and children and their ideas for contextual-appropriate interventions to improve maternal and child nutrition in rural Northern Ghana. Our study revealed that socio-economic factors recognised to influence access to nutritious food



included poverty, poor farm yields and lack of water for dry season agriculture as the main barriers to good nutrition. Subsistence agriculture is the predominant occupation in the study area. Communities depend mainly on rainfall for their farming activities. Not surprisingly, community members mentioned low soil fertility and unreliable rainfall as having negative effects on food production and hence nutritional status of women and children in the area. There were no ethnic differences in views reported on the issues in the current study. It has been previously reported that low agricultural productivity coupled with reliance on rain-fed and low performing largely un-irrigated agricultural lands negatively affects community nutrition⁽⁵⁾. To make matters worse, high levels of poverty in the area make it extremely difficult for farmers to get agricultural inputs such as fertiliser and pesticides to use on their farms in order to boost farm yields. Our findings support earlier studies that reported low income levels as responsible for malnutrition^(26,27). Others have noted that high cost of hiring farm machinery, inadequate access to credit facilities, poor water supply for irrigation and ineffective technical assistance significantly affected nutrition especially in rural communities⁽²⁸⁾.

Socially entrenched gender roles and the lack of fulfillment of responsibilities on the part of the men were also widely discussed, mostly by the women. Generally, men are the head of the household and are supposed to provide for the up-keep of the family. Men's participation and share of their economic responsibilities within the family were considered inadequate and perceived as a barrier to ensuring optimal nutritional status of family members. Although it was the women who voiced the lack of support from the men, some men admitted their lack of involvement⁽¹⁸⁾. Men attributed their inability to provide for the family to the general lack of economic opportunities and the increasingly poor returns from agricultural activities. Also, the excessive drinking habits of some men in the area affected their ability to provide for their family members. Community members observed that lack of financial support coupled with lack of income generating opportunities often made it difficult for families to have access to nutritious food. Evidence exists that men's involvement in child care and feeding could improve nutritional status in rural communities^(29–31).

Despite these numerous challenges, community members proposed various interventions that could improve maternal and child nutrition. They noted that the availability and support to undertake income generating opportunities such as farming and trading especially for women could help solve the poverty situation in the area. They expressed the view that such initiatives would significantly improve maternal and child nutrition largely because the income generated from these activities would make it possible for households to purchase nutritious foods that are not grown in the community. This is consistent with earlier studies

reporting that women's empowerment, employment and social protection could reduce maternal undernutrition, morbidity and mortality^(8,32).

The recommendations from the community members in the current study suggest that a focus on agricultural interventions and women's empowerment would be effective. These types of nutrition-sensitive interventions which also create economic opportunities have been found to improve maternal and child nutrition outcomes⁽³³⁾. Components of interventions which combine agricultural inputs and women's empowerment include training women to set up and manage poultry-based small businesses and upskilling wider community members in home gardening and food demonstrations. Participants in the current study made a point of asking for nutrition education in the form of food demonstrations to enhance their knowledge on the types of locally produced foods they could eat to improve maternal and child nutrition. These types of interventions have been found to significantly improve child dietary diversity and increase infant length, height and weight *z* scores in Ghana⁽⁹⁾. Another agricultural intervention in the neighbouring West African country Burkina Faso also focused on women's empowerment by dedicating land to women and provided training on agriculture production and income generation. This also produced significant reductions in child wasting and improvements in mothers' nutrition and empowerment outcomes^(34,35). Community members in the present study suggested that these types of nutrition-sensitive intervention might be acceptable to them; they wanted support with agricultural outputs, opportunities for women to generate income and education on how to prepare locally grown foods.

Furthermore, measures such as provision of irrigation infrastructure and improved access to agricultural inputs such as high yield seeds, pesticides and fertiliser to boost agricultural productivity were suggested in the current study as ways to improve nutritional status. Study participants suggested that such measures would increase food production and availability all year round and make it easier for households to consume nutritious foods. Evidence from other studies suggests that investment in irrigation infrastructure could be an important poverty alleviation strategy since it boosts agricultural productivity by reducing the risks associated with unreliable rainfall in most of the sub-Saharan African countries including Ghana^(36,37). Small-scale irrigation schemes in the North Eastern region of Ghana have shown to be successful in poverty reduction through creating employment, improving household income source and nutritional status⁽³⁸⁾. Community members spoke of changing climates and unreliable rainfall impacting their agricultural outputs and the food they were able to grow and sell. The 2019 Lancet commission on the global 'syndemic' of obesity, undernutrition and climate change described the way climate change was exacerbating undernutrition



and obesity⁽³⁹⁾. In the context of Navrongo, community members aligned agricultural issues, associated with climate change, to limited access to food and underweight mothers and children. Interventions in this context therefore need to account for effects of climate change and fluctuations in the traditional patterns of the seasons.

Continued investments in nutrition-sensitive interventions could help prevent maternal and child undernutrition, micronutrient deficiencies and mortality especially in rural communities^(8,33). Also, involvement of community members in nutrition interventions could help them identify their own initiatives to improve nutrition^(7,16). It has been demonstrated that good nutrition in early life is the foundation for long-term good health and a healthy dietary pattern reduces maternal and child undernutrition^(40–42). If the proposed strategies identified by study participants are implemented and sustained, this could help to improve maternal and child nutrition in the area.

Strengths and limitations

A strength of this research is the random selection and representation of men and women of different age groups and ethnicity in the FGD. This allowed us to reflect the diversity of experience of men and women in this community and a range of information on potential interventions to improve maternal and child nutrition. The main limitation of the study was that the interviews were conducted in the local languages of the study area, tape recorded, transcribed and translated into English. It is possible that some statements made in the local languages may have lost their original meaning in the English translation. However, the interviews were transcribed by graduate research officers who are native speakers with experience in transcribing qualitative interviews. Any loss of meaning during the translation was minimised and did not affect the findings of the study.

Conclusion

Nutrition plays an important role in maternal and child health. Based on our interpretation of the data, we recommend specific contextual and nutrition-sensitive interventions such as improved irrigation of agricultural land, provision of agricultural inputs to improve crop yield in combination with community education about how to maximise the nutritional benefits of locally available foods as avenues to improve maternal and child nutrition. The impacts of climate change also need to be accounted for, however, when designing agricultural interventions to improve maternal and child nutrition in this context.

It is also recommended that community members' involvement in designing and implementing practical nutrition programmes at the community level could lead to

improvement in maternal and child nutritional intake and sustainability of such programmes. It is therefore important for stakeholders such as the Ministry of Food and Agriculture, Ministry of Health, Ghana Health Service and civil society organisations to take appropriate steps towards supporting community initiated strategies to improving maternal and child nutrition in Ghana.

Acknowledgements

Acknowledgements: The authors would like to express their profound gratitude to all the study participants for sharing their experiences and views with the research team. We are also grateful to all individuals who supported the study team in one way or the other during data collection and analysis. *Financial support:* The current research was funded by the National Institute for Health Research (NIHR) (17\63\154) using UK aid from the UK Government to support global health research. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care. *Conflict of interest:* There are no conflicts of interest. *Authorship:* Substantial contributions to conception and design: C.D., E.A.N., M.-L.N. and M.B. Data acquisition: C.D., J.K.A., E.D., P.B. and E.W.N. Analysis and/or interpretation: C.D., S.T.C., J.K.A., E.D., R.A. and M.B. Drafting the article: C.D., S.T.C., J.K.A., E.W.N. and E.D. Critically revised the article for important intellectual content: E.A.N., D.W., S.H.K., M.A.D., W.O., R.A., P.W., A.R.O., M.-L.N. and M.B. Final approval of the article: C.D., E.A.N., S.T.C., J.K.A., E.D., P.B., E.W.N., D.A.-B., D.W., S.H.K., M.A.D., W.O., R.A., P.W., A.R.O., M.-L.N. and M.B. *Ethics of human subject participation:* The current study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the Navrongo Health Research Centre Institutional Review Board (approval no. NHRCIRB322) and the University of Southampton Faculty of Medicine Ethics Committee (no. 47290). Written informed consent was obtained from all subjects/patients.

References

1. Bhutta ZA, Ahmed T, Black RE *et al.* (2008) What works? Interventions for maternal and child undernutrition and survival. *Lancet* **371**, 417–440.
2. Ghana Statistical Service, Ghana Health Service & ICF International (2015) *Ghana Demographic and Health Survey 2014*. Rockville, MD: GSS, GHS and ICF International; available at <https://dhsprogram.com/pubs/pdf/FR307/FR307.pdf> (accessed June 2019).
3. Sienso G & Lyford C (2018) Assessing the factors affecting malnutrition in northern Ghana. *J Nutr Disorders Ther* **8**, 235.
4. Doku DT & Neupane S (2015) Double burden of malnutrition: increasing overweight and obesity and stall underweight



- trends among Ghanaian women. *BMC Public Health* **1**, 670.
5. Food and Agricultural Organization (2015) Country fact sheet on food and agriculture policy trends; available at <http://www.fao.org/3/a-i4490e.pdf> (accessed June 2020).
 6. National Nutrition Policy (2013) National Nutrition Policy for Ghana, 2013–2017; available at <http://extwprlegs1.fao.org/docs/pdf/gha145267.pdf> (accessed June 2020).
 7. Masters WA, Rosettie K, Kranz S *et al.* (2017) Priority interventions to improve maternal and child diets in sub-Saharan Africa and South Asia. *Matern Child Nutr* **14**, e12526.
 8. Bhutta ZA, Das JK, Rizvi A *et al.* (2013) Maternal and child nutrition 2: evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet* **382**, 452–477.
 9. Marquis GS, Colecraft EK, Kanlisi R *et al.* (2018) An agriculture–nutrition intervention improved children’s diet and growth in a randomized trial in Ghana. *Matern Child Nutr* **14**, e12677.
 10. Malapit HJ & Quisumbing AR (2015) What dimensions of women’s empowerment in agriculture matter for nutrition in Ghana? *Food Policy* **1**, 54–63.
 11. Gongwer CR & Aryeetey R (2014) Implementing nutrition interventions in Ghana at district level: gaps and opportunities. *Afr J Food Agric Nut Dev (AIFAND)* **14**, 1684–5374.
 12. Zakari S, Ying L & Song B (2014) Factors influencing household food security in West Africa: the case of southern Niger. *Sustainability* **6**, 1191–1202.
 13. Etwire PM, Al-Hassan RM, Kuwornu JK *et al.* (2013) Application of livelihood vulnerability index in assessing vulnerability to climate change and variability in Northern Ghana. *J Environ Earth Sci* **3**, 157–170.
 14. Antwi-Agyei P, Fraser ED, Dougill AJ *et al.* (2012) Mapping the vulnerability of crop production to drought in Ghana using rainfall, yield and socioeconomic data. *Appl Geogr* **1**, 324–334.
 15. Tette EMA, Sifah EK & Nartey ET (2015) Factors affecting malnutrition in children and the uptake of interventions to prevent the condition. *BMC Pediatr* **15**, 189.
 16. Robinson SM (2018) Improving nutrition to support healthy ageing: what are the opportunities for intervention? *Proc Nutr Soc* **77**, 257–264.
 17. Naidu S, Baliga SS, Yadav H *et al.* (2017) Psychological factors affecting dietary habits of college going adolescents (17–19 years) in urban area of Belgaum. *Intl J Med Sci Public Health* **6**, 67–70.
 18. Compaoré A, Ouedraogo K, Boua R *et al.* (2020) “Men are no more playing their roles”: maternal and child nutrition in Nanoro, Burkina Faso. *Public Health Nutr* (unpublished).
 19. Erzse A, Goldstein S, Norris SA *et al.* (2020) Double-duty solutions for optimising maternal and child nutrition in urban South Africa: a qualitative study. *Public Health Nutr* (unpublished).
 20. Watson D, Kehoe SH, Erzse A *et al.* (2020) Community perspectives of maternal and child health during nutrition and economic transition in sub-Saharan Africa. *Public Health Nutr* (unpublished).
 21. Creswell JW (2009) *Research Design: Qualitative, Quantitative, and Mixed Methods Approaches*, 3rd ed. Thousand Oaks CA: Sage Publications, 260.
 22. Oduro AR, Wak G, Azongo D *et al.* (2012) Profile of the Navrongo Health and Demographic Surveillance System. *Int J Epidemiol* **41**, 968–976.
 23. Chatio S & Akweongo P (2017) Retention and sustainability of community-based health volunteers’ activities: a qualitative study in rural Northern Ghana. *PLoS ONE* **12**, e0174002.
 24. Chatio S, Aborigo R, Adongo PB *et al.* (2016) Factors influencing adverse events reporting within the health care system: the case of artemisinin-based combination treatments in northern Ghana. *Malar J* **15**, 125.
 25. Guest G, Macqueen K & Namey EE (2012) *Applied Thematic Analysis*. Thousand Oaks, CA: SAGE Publications, Inc.
 26. Yimer M (2018) Review on factors affecting the nutritional status of women in Ethiopia. *Interventions Obes Diabetes* **2**, IOD.000526.2018.
 27. Panth N, Gavarkovs A, Tamez M *et al.* (2018) The influence of diet on fertility and the implications for public health nutrition in the United States. *Front Public Health* **6**, 211.
 28. Armah S (2016) Strategies to stimulate Ghana’s economic transformation and diversification. *Ashesi Econ Lect Ser J* **2**, 9–16.
 29. Juma OA, Enumah ZO, Wheatley H *et al.* (2016) Prevalence and assessment of malnutrition among children attending the Reproductive and Child Health clinic at Bagamoyo District Hospital, Tanzania. *BMC Public Health* **16**, 1094.
 30. Kansime N, Atwine D, Nuwamanya S *et al.* (2017) Effect of male involvement on the nutritional status of children less than 5 years: a cross sectional study in a rural southwestern district of Uganda. *J Nutr Metab* **9**, 3427087.
 31. Abate KH & Belachew T (2017) Women’s autonomy and men’s involvement in child care and feeding as predictors of infant and young child anthropometric indices in coffee farming households of Jimma Zone, South West of Ethiopia. *PLoS ONE* **12**, e0172885.
 32. Rahman MM, Saima U & Goni MA (2015) Impact of maternal household decision-making autonomy on child nutritional status in Bangladesh. *Asia-Pacific J Public Health* **27**, 509–520.
 33. Ruel MT & Alderman H (2013) Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet* **382**, 536–551.
 34. Heckert J, Olney DK & Ruel MT (2019) Is women’s empowerment a pathway to improving child nutrition outcomes in a nutrition-sensitive agriculture program? Evidence from a randomized controlled trial in Burkina Faso. *Soc Sci Med* **233**, 93–102.
 35. Olney DK, Pedehombga A, Ruel MT *et al.* (2015) A 2-year integrated agriculture and nutrition and health behavior change communication program targeted to women in Burkina Faso reduces anemia, wasting, and diarrhea in children 3–12.9 months of age at baseline: a cluster-randomized controlled trial. *J Nutr* **145**, 1317–1324.
 36. Dinye RD (2013) Irrigated agriculture and poverty reduction in Kassena-Nankana district in the upper east region, Ghana. *J Sci Tech* **33**, 59–72.
 37. Hoddinott J, Berhane G, Gilligan DO *et al.* (2012) The impact of Ethiopia’s productive safety net programme and related transfers on agricultural productivity. *J Afr Econ* **21**, 761–786.
 38. Adam JN, Al-hassan S & Akolgo DA (2016) Small scale irrigation and rural poverty reduction in the upper east region of Ghana. *Afr J Sci Res* **5**, 2.
 39. Swinburn BA, Kraak VI, Allender S *et al.* (2019) The global syndemic of obesity, undernutrition, and climate change: the Lancet Commission report. *Lancet* **393**, 791–846.
 40. Mohammed YHI, Khanum NF, Mamatha SV *et al.* (2016) The prevalence of malnutrition in developing countries: a review. *J Bio Innov* **5**, 810–815.
 41. Cetin I & Laoreti A (2015) The importance of maternal nutrition for health. *J Pediatr Neonatal Individualized Med* **4**, e040220.
 42. Kumar K (2017) Importance of healthy life style in healthy living. *JOJ Pub Health* **2**, 555596.