

# Technical note

## Unidentified specimens in zooarchaeology

Shaw Badenhorst<sup>1,2\*</sup> & Ina Plug<sup>3</sup>

<sup>1</sup>Archaeozoology Section, Ditsong National Museum of Natural History  
(former Transvaal Museum),  
P.O. Box 413, Pretoria, 0001 South Africa

<sup>2</sup>Institute for Human Evolution, University of the Witwatersrand,  
Private Bag 3, Wits, 2050 South Africa

<sup>3</sup>Department of Anthropology and Archaeology, University of South Africa,  
c/o P.O. Box 21022, Valhalla, 0137 South Africa

Received 25 July 2011. Accepted 25 October 2011

### INTRODUCTION

A consideration of skeletal part profiles of faunal assemblages has become more or less part of conventional zooarchaeological analyses throughout the world. Studies of skeletal parts yield a wealth of information, and can inform zooarchaeologists on, for example, site function and usage (Binford 1978), carcass transportation (Marean *et al.* 1992), meat provisioning, sumptuary rules and social status (Ijzereef 1989; Schmitt & Lupo 2008), gender (Mooketsi 2001), feasting (Hayden 2001), trade or long-distance meat acquisition (Driver 1990), bone preservation (Brain 1967), as well as attributes of non-human agents such as carnivores (Marean & Spencer 1991), raptors and porcupines (Brain 1981).

The vast majority of faunal assemblages from Pleistocene and Holocene deposits in all parts of the world, regardless of the agent(s) of accumulation, consist of fragmented specimens. The fragmentary nature of these faunal assemblages is due to a complex interaction between biotic and abiotic processes, which transform complete or near-complete elements into fragments over time. These taphonomic processes include, for example, cutting, chopping, burning, cooking, trampling, weathering, rootlet etching, carnivore and rodent gnawing, digestion, diagenetic actions and retrieval damage (e.g. Lyman 1994; Reitz & Wing 1999). Complete elements are usually associated with animal burials (Muir & Driver 2004, but see Hutten 2008), which are not considered here. An aspect most often ignored in studies of animal skeletal profiles in zooarchaeology, is unidentified specimens.

### UNIDENTIFIED SPECIMENS

The first step in faunal analysis is often to separate identifiable from unidentifiable specimens. In most cases, a large percentage of specimens in a faunal assemblage cannot be identified (O'Connor 2000; Table 1). The ratio of identified and unidentified specimens provides some indication of the level of fragmentation in an assemblage (Plug 1988), although this is also influenced by the method of analyses. For example, the method suggested

by Driver (2005) regards all specimens that can be identified to a skeletal element as 'identifiable', whereas Brain (1974) and Voigt (1983) regard indeterminate enamel fragments, skull, vertebrae and ribs as 'unidentified' specimens.

In addition, archaeological recovery methods are also important. A lack of or poor screening and handpicking of specimens during excavations can inflate the ratio of identified to unidentified specimens. For example, the faunal assemblage from the Late Iron Age site of Simunye in Swaziland, comprised 62% identified specimens (Badenhorst & Plug 2002), which was due to a lack of screening during excavations (compare Table 1). Objects made from bone and shell are not always presented to the faunal analyst and where such objects are common, for example beads, bone points, tortoise shell containers and pendants, their absence from the faunal sample to be analysed may cause misrepresentation of taxa and skeletal elements. The proficiency of zooarchaeologists is also an important factor. In this regard, O'Connor (2000) points out that 'unidentified' specimens are not the same as those that are 'unidentifiable'. The former category indicates that the zooarchaeologist did not regard that sufficient morphological criteria were present to allow taxonomic identification. This O'Connor (2000: 42) correctly terms a positive decision, not an admission of defeat. In fact, some specimens that may be considered 'unidentified' by one zooarchaeologist is often not for another (Grayson 1984). Controversially, Binford & Bertram (1977: 125) state that, in referring to both archaeological and ethnographic faunal assemblages, they '...have always taken the position that there is no unidentifiable bone. All bones, even the smallest fragments, may be identified with sufficient training in osteology' (emphasis by original authors). Such statements O'Connor (2000: 42) correctly calls '...wonderfully optimistic, and utterly wrong.'

The nature of an assemblage may also determine the ratio of unidentifiable to identifiable specimens. When a collection contains a large amount of fragments that are imminently identifiable such as tortoise shell for example, and the same sample also contains a large number of fragmented mammal bones that cannot be identified, it may lead to problems of interpretation. The identifiability of certain groups is demonstrated by the assemblage from Likoeng, a Later Stone Age site in Lesotho that was occupied between c. 4000 and 1200 BP. The faunal assemblage consists of c. 1 680 000 specimens of which c. 1.3 million specimens (77%) are fish remains (Plug & Mitchell 2008). Of the fish remains, over 55% could be identified to taxon or genus, while for the non-fish remains, less than 2% could be identified to taxon or genus (Plug 2006, unpubl. data).

Bone breakage tends to, initially at least increase the percentage of Number of Identified Specimens (NISP) in an assemblage. However, when an assemblage is heavily fragmented the opposite effect is produced. A greater proportion of specimens then become too small to be identified, and hence decrease the percentage of identified specimens (Marshall & Pilgram 1993). Lyman &

\*Author for correspondence. E-mail: shaw@ditsong.org.za

**Table 1.** Examples of faunal assemblages from southern Africa indicating level of fragmentation (percentages rounded off).

Sites	Date and cultural association	% Identified specimens (NISP)	% Unidentified specimens	Reference
Steinaecker's Horse	AD 1899–1902 (Historical Period)	27	73	Badenhorst <i>et al.</i> 2002
Boleu	Mid AD 1800s (Terminal Late Iron Age)	17	83	Badenhorst & Plug 2004–2005
Boitsemagano	17th century (Late Iron Age)	13	87	Plug & Badenhorst 2006
Manyikeni	12th–16th/17th century (Late Iron Age)	27	73	Sigvallius 1988
Ratho	AD 1040–1240 (Middle Iron Age)	14	86	Brunton 2010
KwaGandaganda	AD 620–1030 (Early Iron Age)	23	77	Beukes 2000
Toteng 1	c. 2070–1480 BP (Pastoral Later Stone Age)	11	89	Robbins <i>et al.</i> 2008, Badenhorst, unpubl. data
iNkolimahashi Shelter (excluding rodents)	c. 360–3130 BP (Later Stone Age)	8	92	Badenhorst 2003
Olieboomspoor	<2000 BP (Later Stone Age)	33	67	Van der Ryst 2006
Maqonqo Shelter	3500–9000 BP (Later Stone Age)	2	98	Plug 1996
Sibudu Cave	Pre-38 000 BP (Middle Stone Age)	7	93	Plug 2004
Florisbad	Middle–Late Pleistocene (Middle Stone Age)	45	55	Brink 1987

O'Brien (1987: 496) pointed out that when specimens are '...reduced beyond the minimal identifiable size, then the proportion of identifiable fragments will be decreased.' It can therefore happen that, in samples with relatively high percentages of unidentified specimens, most of the absent skeleton was deposited, and in fact, recovered during excavations. However, the high percentage of unidentified specimens can preclude species or element identification (also Todd & Rapson 1988; Watson 1972). Lyman & O'Brien (1987: 496) described this effect as an 'analytic absence' of species and elements.

### CASE STUDIES

We present three examples of how a consideration of unidentified specimens can lead zooarchaeologists to different interpretations. At the Middle Stone Age site of Klasies River Mouth, Klein (1976, 1989) found that the upper limb bones of larger bovids are generally absent, whereas lower limb bones and the head are common. In contrast, smaller bovids are represented by more even skeletal part representation. The 'schlepp effect' featured strongly to explain this pattern, whereby hunters discarded the bulkier limb bones of large animals at kill-sites and dragged the meat back to camp in the skin using the intact foot bones as handles (Perkins & Daly 1968). In a reply, Bartram & Marean (1999) showed that unless unidentified long bone shafts are refitted to form more complete bones, the upper limbs of large animals will be under-represented.

However, refitting is tedious and time-consuming (Klein *et al.* 1999). In addition, some faunal assemblages such as those from the Middle Stone Age are often reduced to crumbs. For example, at the Middle Stone Age site of Sibudu Cave, the vast majority of bone remains were reduced to small pieces, making it even impossible to recognize long bone flakes (*cf.* Plug 2004). Nonetheless, the research by Bartram & Marean (1999) highlighted the importance of unidentified long bones.

Reynard (2011, also J. Reynard, S. Badenhorst & C.S. Henshilwood, in prep.) studied the unidentified long

bones from the Middle Stone Age layers of Blombos Cave. The identified remains (Henshilwood *et al.* 2001) indicate that small animals such as rock hyrax, Cape dune molerat and small bovids dominate the faunal sample. However, the cortical thickness of the unidentified long bones indicates that medium-sized animals were more common than smaller game. However, issues related to identification and taphonomy may also have caused this pattern.

Studies of faunal assemblages from Middle Period and early Plateau Pithouse Tradition (7000–3500 BP) sites on the Interior Plateau of British Columbia, Canada, yielded a dominance of medium (deer-sized) and large (elk-sized) artiodactyla lower limb bones. Although this pattern is consistent with results of ethnoarchaeological studies of butchering camps (Binford 1978), the very high percentage of unidentified, and very low percentage of identified specimens (Table 2), suggests that more, and missing elements of artiodactyls are probably present in assemblages, but these could not be identified. Most of the assemblage has been reduced to crumbs (Badenhorst 2009). Although it remains likely that these sites were butchering camps, the skeletal profile is more likely a taphonomic artefact rather than conforming to an ethnoarchaeological pattern (Binford 1978).

### CONCLUSIONS

Skeletal part profiles form a central part understanding bone taphonomy. A factor most often overlooked when evaluating skeletal part profiles, is unidentified specimens. Faunal analysts must be aware that any profile of skeletal parts reflects a pattern within the *identified* sample, and not necessarily within the *entire* assemblage, especially in assemblages with high percentages of unidentified specimens. Conversely, in assemblages with high percentages of identified, and low percentages of unidentified specimens, any pattern of skeletal part profiles may be assumed to reflect the entire assemblage. However, it is as yet impossible to determine what constitute 'high percentages' of unidentified specimens. Ethnoarchaeological, actualistic and modelling studies may provide greater

**Table 2.** Middle Period and early Plateau Pithouse Tradition samples (Badenhorst 2009).

Site	% Identified specimens (NISP)	% Unidentified specimens
EdRh-31	6	94
EeRh-61	10	90
EdRh-11	1	99
EdRh-14	3	97

insights. It is useful when such studies list the number of unidentified specimens (e.g. Lupo 2001).

The role of unidentified specimens in skeletal part profiles will likely remain a problematic issue. Each assemblage has a unique taphonomic history. Nonetheless, zooarchaeologists can make a contribution by:

- providing an explicit indication about the method used to analyse an assemblage (Driver 1982);
- presenting the numbers and percentages of identified and unidentified specimens for assemblages (Grigson 1978); and
- in cases where the method of Brain (1974) and Voigt (1983) is used, presenting the numbers of unidentified fragments that could not be placed into a taxonomic category, under enamel, skull, vertebrae and rib.

In addition, it is important that archaeologists retain all excavated bone specimens, and subject all of these for analysis. Most bone specimens can be retained using a 1-mm sieve, but in some instances the use of smaller sieve sizes may be advisable (Matsui 2008). Depending on the research questions, preservation and resources, refitting may be a viable option in some cases. By being more explicit about the analytical method and ratio of identifiable *versus* unidentified specimens, zooarchaeologists will be in a better position to evaluate interpretations based on skeletal part profiles. While it is tempting to immediately associate any changes in skeletal part profiles to human behaviour, such patterns could also relate to changes in the ratio of identified *versus* unidentified specimens.

However, many studies have applied skeletal part profiles successfully. For example, Driver (1990) noted that those bison elements most likely affected by post-depositional destruction – ribs – dominate faunal samples in the Sierra Blanca region of southeastern New Mexico for sites dating to between AD 1150 and 1450. Taphonomic processes were largely excluded, as more dense elements such as distal humeri, phalanges and distal tibiae are absent or occur in very low frequencies. In addition, ethnographies from the region indicate that brisket meat cuts are highly prized (Driver 1990).

Many archaeologists and zooarchaeologists may not find a consideration of unidentified specimens, and its potential effects on interpretations appealing (but see Badenhorst 2009; Thackeray 2007; Reynard 2011). In this regard, Maltby (2002: 88–89) remarks: ‘There is a widely held belief that animal bone studies have failed to produce the answers to what other archaeologists want to know. For example, when an archaeozoologist is asked

the apparent simple question of ‘What did they eat?’, there is likely to be a very convoluted answer. Of course, archaeozoologists are right to point out the complexities caused by taphonomic processes; small sample size; retrieval rates; intra- and inter-site variation; the effects of different methods of quantification, etc. However, it’s a sad fact that others are not particularly interested in such problems. They want positive answers. Counter-arguments to the effect that at least archaeozoologists are attempting to look at their data critically, although valid, are unpopular.’

It may be worth pointing out that the aims and purpose of archaeology is to understand the past, and how humans interacted with each other and the animals, environment and material culture (*cf.* Brinton 1895; Petrie 1904). At the same time, it is imperative to realize the limitations of our data, and that the archaeological record is ‘...the imperishable remnants of material culture, not the sum of artefacts in use by some particular people at some particular time’ (Summers 1958: 6).

The two reviewers offered constructive suggestions, for which we are grateful.

## REFERENCES

- BADENHORST, S. 2009. Artiodactyla skeletal parts at Middle Period and early Plateau Pithouse Tradition sites on the Interior Plateau, British Columbia: a view from EdRh-31. *Canadian Zooarchaeology* **26**, 25–41.
- BADENHORST, S. 2003. The archaeofauna from iNkolimahashi Shelter, a Later Stone Age shelter in the Thukela Basin, KwaZulu-Natal, South Africa. *Southern African Humanities* **15**, 45–57.
- BADENHORST, S. & PLUG, I. 2002. Appendix: Animal remains from recent excavations at a Late Iron Age site, Simunye, Swaziland. *Southern African Humanities* **14**, 45–50.
- BADENHORST, S., PLUG, I., PELSER, A.J. & VAN VOLLENHOVEN, A.C. 2002. Faunal analysis from Steinaecker’s Horse, the northernmost British military outpost in the Kruger National Park during the South African War. *Annals of the Transvaal Museum* **39**, 57–63.
- BADENHORST, S. & PLUG, I. 2004–2005. Boleu: faunal analysis from a 19th century site in the Groblersdal area, Mpumalanga, South Africa. *Southern African Field Archaeology* **13&14**, 13–18.
- BARTRAM, L.E. & MAREAN, C.W. 1999. Explaining the ‘Klasies Pattern’: Kwa ethnoarchaeology, the Die Kelders Middle Stone Age archaeofauna, long bone fragmentation and carnivore ravaging. *Journal of Archaeological Science* **26**, 9–12.
- BEUKES, C.F. 2000. *KwaGandaganda: an archaeozoological case study of the exploitation of animal resources during the Early Iron Age in KwaZulu-Natal*. M.A. thesis. Pretoria, University of South Africa.
- BINFORD, L.R. 1978. *Numamiut ethnoarchaeology*. London, Academic Press.
- BINFORD, L.R. & BERTRAM, J.B. 1977. Bone frequencies—and attritional processes. In: Binford, L.R., (ed.), *For Theory Building in Archaeology. Essays on Faunal Remains, Aquatic Resources, Spatial Analysis, and Systematic Modeling*, 77–153. London, Academic Press.
- BRAIN, C.K. 1967. Hottentot food remains and their bearing on the interpretation of fossil bone assemblages. *Scientific Papers of the Namib Desert Research Station* **32**, 1–7.
- BRAIN, C.K. 1974. Some suggested procedures in the analysis of bone accumulations from southern African Quaternary sites. *Annals of the Transvaal Museum* **29**, 1–8.
- BRAIN, C.K. 1981. *The Hunters or the Hunted? An Introduction to African Cave Taphonomy*. Chicago, University of Chicago Press.
- BRINK, J.S. 1987. The archaeozoology of Florisbad, Orange Free State. *Memoirs van die Nasionale Museum Bloemfontein* **24**. Bloemfontein, National Museum.
- BRINTON, D.G. 1895. The aims of anthropology. *Science (New Series)* **2(35)**, 241–252.
- BRUNTON, S. 2010. *Ritual fauna from Ratho Kroonkop: an early second millennium AD rain-control site in the Shashe Limpopo Confluence Area, South Africa*. Honours thesis. Pretoria, University of Pretoria.
- DRIVER, J.C. 1982. Minimum standards for reporting of animal bones in salvage archaeology: southern Alberta as a case study. In: Francis, P.D. & Poplin, E.C. (eds), *Directions in Archaeology. A Question of Goals*,

- 199–209. Calgary, The Archaeological Association of the University of Calgary.
- DRIVER, J.C. 1990. Bison assemblages from the Sierra Blanca region, southeastern New Mexico. *Kiva* 55(3), 245–263.
- DRIVER, J.C. 2005. Manual for description of vertebrate remains. Unpublished manual. Cortez, Crow Canyon Archaeological Center.
- GRAYSON, D.K. 1984. *Quantitative Zooarchaeology. Topics in the Analysis of Archaeological Faunas*. London, Academic Press.
- GRIGSON, C. 1978. Towards a blueprint for animal bone reports in archaeology. In: Brothwell, D.R., Thomas, K.D. & Clutton-Brock, J. (eds), *Research Problems in Zooarchaeology*, 121–128. Occasional Paper No. 3, Institute of Archaeology. London, University of London.
- HAYDEN, B. 2001. Fabulous feasts. A prolegomenon to the importance of feasting. In: Dietler, M. & Hayden, B., (eds), *Feasts. Archaeological and Ethnographic Perspectives on Food, Politics, and Power*, 23–64. Washington, Smithsonian Institution Press.
- HENSHILWOOD, C.S., SEALY, J.C., YATES, R., CRUZ-URIBE, K., GOLDBERG, P., GRINE, F.E., KLEIN, R.G., POGGENPOEL, C., VAN NIEKERK, K. & WATTS, I. 2001. Blombos Cave, Southern Cape, South Africa: preliminary report on the 1992–1999 excavations of the Middle Stone Age levels. *Journal of Archaeological Science* 28, 421–448.
- HUTTEN, L. 2008. Symbolic animal burials from the Venda region in the Limpopo Province, South Africa. In: Badenhorst, S., Mitchell, P. & Driver, J.C. (eds), *Animals and People: Archaeozoological Papers in Honour of Ina Plug*, 186–199. Oxford, British Archaeological Reports 1849.
- IJZEREEF, F.G. 1989. Social differentiation from animal bone studies. In: Serjeantson, D. & Waldron, T. (eds), *Diets and Crafts in Towns*, 42–53. Oxford, British Archaeological Reports 199.
- KLEIN, R.G. 1976. The mammalian fauna of the Klasies River Mouth sites, Cape Province, South Africa. *South African Archaeological Bulletin* 31, 75–98.
- KLEIN, R.G. 1989. Why does skeletal part representation differ between smaller and larger bovids at Klasies River Mouth and other archaeological sites? *Journal of Archaeological Science* 6, 363–381.
- KLEIN, R.G., CRUZ-URIBE, K. & MILO, R.G. 1999. Skeletal part representation in archaeofaunas: comments on 'Explaining the 'Klasies Pattern': Kua ethnoarchaeology, the Die Kelders Middle Stone Age archaeofauna, long bone fragmentation and carnivore raving' by Bartram & Marean. *Journal of Archaeological Science* 26, 1225–1234.
- LUPO, K.D. 2001. Archaeological skeletal part profiles and differential transport: an ethnoarchaeological example from Hadza bone assemblages. *Journal of Anthropological Archaeology* 20, 361–378.
- LYMAN, R.L. 1994. *Vertebrate Taphonomy*. Cambridge, Cambridge University Press.
- LYMAN, R.L. & O'BRIEN, M.J. 1987. Plow-zone zooarchaeology: fragmentation and identifiability. *Journal of Field Archaeology* 14(4), 493–498.
- MALTBY, M. 2002. Animal bones in archaeology: how archaeozoologists can make a greater contribution to British Iron Age and Romano-British archaeology. In: Dobney, K. & O'Connor, T. (eds), *Bones and the Man. Studies in Honour of Don Brothwell*, 88–94. Oxford, Oxbow Books.
- MAREAN, C.W. & SPENCER, L.M. 1991. Impact of carnivore ravaging on zooarchaeological measures of element abundance. *American Antiquity* 56, 645–658.
- MAREAN, C.W., SPENCER, L.M., BLUMENSCHINE, R.J. & CAPALDO, S.D. 1992. Captive hyaena bone choice and destruction, the Schlepp Effect, and Olduvai archaeofaunas. *Journal of Archaeological Science* 19, 101–121.
- MARSHALL, F. & PILGRAM, T. 1993. NISP vs MNI in quantification of body-part representation. *American Antiquity* 58(2), 261–269.
- MATSUI, A. 2008. *Fundamentals of Zooarchaeology in Japan*. Kyoto, Kyoto University Press.
- MOOKETSI, C. 2001. Butchering styles and the processing of cattle carcasses in Botswana. *Pula: Botswana Journal of African Studies* 15(1), 108–124.
- MUIR, R.J. & DRIVER, J.C. 2004. Identifying ritual use of animals in the northern American Southwest. In: O'Day, S.J., Van Neer, W. & Eryvnyck, A. (eds), *Behaviour Behind Bones. The Zooarchaeology of Ritual, Religion, Status and Identity*, 128–143. Oxford, Oxbow Books.
- O'CONNOR, T. 2000. *The Archaeology of Animal Bones*. College Station, Texas A&M University Press.
- PERKINS, D. & DALY, P. 1968. A hunters' village in Neolithic Turkey. *Scientific American* 219, 96–106.
- PETRIE, W.M.F. 1904. *Methods and Aims in Archaeology*. London, Macmillan.
- PLUG, I. 1988. *Hunters and herders: an archaeozoological study of some prehistoric communities in the Kruger National Park*. D.Phil. dissertation. Pretoria, University of Pretoria.
- PLUG, I. 1996. The hunter's choice: faunal remains from Maqonqo Shelter, South Africa. *Natal Museum Journal of Humanities* 8, 41–52.
- PLUG, I. 2004. Resource exploitation: animal use during the Middle Stone Age at Sibudu Cave, KwaZulu-Natal. *South African Journal of Science* 100, 151–158.
- PLUG, I. 2006. The exploitation of freshwater fish during the Later Stone Age of Lesotho. In: Grier, C., Kim, J. & Uchiyama, J. (eds), *Beyond Affluent Foragers: Rethinking Hunter-gatherer Complexity*, 24–33. Oxford, Oxbow Books.
- PLUG, I. & BADENHORST, S. 2006. Notes on the fauna from three Late Iron Age mega-sites, Boitsemagano, Molokwane and Mabjanamatshwana, North West Province, South Africa. *South African Archaeological Bulletin* 61(183), 57–67.
- PLUG, I. & MITCHELL, P. 2008. Fishing in the Lesotho highlands: 26 000 years of fish exploitation; with special reference to Sehonghong Shelter. *Journal of African Archaeology* 6(1), 33–55.
- REITZ, E.J. & WING, E.S. 1999. *Zooarchaeology*. Cambridge Manuals in Archaeology. Cambridge, Cambridge University Press.
- REYNARD, J. 2011. *The unidentified long bone fragments from the Middle Stone Age Still Bay layers at Blombos Cave, Southern Cape, South Africa*. M.Sc. thesis. Johannesburg, University of the Witwatersrand.
- ROBBINS, L.H., CAMPBELL, A.C., MURPHY, M.L., BROOK, G.A., LIANG, F., SKAGGS, S.A., SRIVASTAVA, P., MABUSE, A.A. & BADENHORST, S. 2008. Recent archaeological and paleoenvironmental research at Toteng, Botswana: early domesticated livestock in the Kalahari. *Journal of African Archaeology* 6(1), 131–149.
- SCHMITT, D.N. & LUPO, K.D. 2008. Do faunal remains reflect socioeconomic status? An ethnoarchaeological study among Central African farmers in the northern Congo Basin. *Journal of Anthropological Archaeology* 27, 315–325.
- SIGVALLIUS, B. 1988. The faunal remains from Manyikeni. In: Sinclair, P.J., Törnblom, M., Bohm, C., Sigvallius, B. & Hulthén, B. (eds), *Analyses of Slag, Iron, Ceramics and Animal Bones from Excavations in Mozambique*, 23–34. Studies in African Archaeology 2. Maputo and Stockholm: Eduardo Mondlane University and the Central Board of National Antiquities.
- SUMMERS, R. 1958. Methods and aims in South African archaeology: Presidential Address, 1957, read 11 March 1958 at Bulawayo. *South African Archaeological Bulletin* 13(49), 3–9.
- THACKERAY, J.F. 2007. Hominids and carnivores at Kromdraai and other Quaternary sites in southern Africa. In: Pickering, T.R., Schick, K. & Toth, N. (eds), *Breathing Life into Fossils: Taphonomic Studies in Honor of C.K. (Bob) Brain*, 44–49. Bloomington, Stone Age Institute Press.
- TODD, L.C. & RAPSON, D.J. 1988. Long bone fragmentation and interpretation of faunal assemblages: approaches to comparative analysis. *Journal of Archaeological Science* 15, 307–325.
- VAN DER RYST, M.M. 2006. *Seeking shelter: Later Stone Age hunters, gatherers and fishers of Olieboomspoor in the western Waterberg, south of the Limpopo*. Ph.D. dissertation. Johannesburg, University of the Witwatersrand.
- VOIGT, E.A. 1983. *Mapungubwe: an Archaeozoological Interpretation of an Iron Age Community*. Transvaal Museum Monograph No. 1. Pretoria, Transvaal Museum.
- WATSON, J.P.N. 1972. Fragmentation analysis of animal bone samples from archaeological sites. *Archaeometry* 14, 221–228.