

Blood flow rate and wall shear stress in seven major cephalic arteries of humans

Roger S. Seymour,¹ Qiaohui Hu¹ and Edward P. Snelling^{2,3}

¹School of Biological Sciences, Faculty of Sciences, University of Adelaide, Adelaide, SA, Australia

²Department of Anatomy and Physiology, Faculty of Veterinary Science, University of Pretoria, Pretoria, South Africa

³Brain Function Research Group, School of Physiology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Abstract

Blood flow rate (\dot{Q}) in relation to arterial lumen radius (r_i) is commonly modelled according to theoretical equations and paradigms, including Murray's Law ($\dot{Q} \propto r_i^3$) and da Vinci's Rule ($\dot{Q} \propto r_i^2$). Wall shear stress (τ) is independent of r_i with Murray's Law ($\tau \propto r_i^0$) and decreases with da Vinci's Rule ($\tau \propto r_i^{-1}$). These paradigms are tested empirically with a meta-analysis of the relationships between \dot{Q} and r_i in seven major arteries of the human cephalic circulation from 19 imaging studies in which both variables were presented. The analysis shows that $\dot{Q} \propto r_i^{2.16}$ and $\tau \propto r_i^{-1.02}$, more consistent with da Vinci's Rule than Murray's Law. This meta-analysis provides standard values for \dot{Q} , r_i and τ in the human cephalic arteries that may be a useful baseline in future investigations. On average, the paired internal carotid arteries supply 75%, and the vertebral arteries supply 25%, of total brain blood flow. The internal carotid arteries contribute blood entirely to the anterior and middle cerebral arteries and also partly to the posterior cerebral arteries via the posterior communicating arteries of the circle of Willis. On average, the internal carotid arteries provide 88% of the blood flow to the cerebrum and the vertebral arteries only 12%.

Key words: artery; brain; cardiovascular; cerebral; da Vinci's Rule; Murray's Law.

Introduction

The structure of the arterial tree has attracted the attention of scholars for centuries. Even before the blood was shown to circulate by Harvey in 1628 (Willis, 2016), Leonardo da Vinci described the branching of the arteries in mathematical terms, indicating that the cross-sectional area of a parent artery was equal to the sum of the cross-sectional areas of the two child arteries branching from it (Richter, 1970), a relationship that is now called 'da Vinci's Rule'. In other words, the square of the diameter of a parent artery should be equal to the sum of the squares of the diameters of the two child arteries. More than three centuries later, in an attempt to analyse the structure of the arterial tree as an efficient energy-conserving system, Cecil Murray proposed that, to minimise energy losses, the cube of the diameter of the parent artery should be equal to the sum of the cubes of the diameters of the two child

arteries, a relationship that became known as 'Murray's Law' (Murray, 1926). Murray's Law was a formal treatment of what was proposed by Thomas Young in 1808, that is, the ratio of the diameters of the parent and child vessels is $2^{1/3} : 1$ (Sherman, 1981). Mathematically, these paradigms can be expressed in terms of vessel radius, and represent two cases of the general relationship:

$$r_p^n = r_{c1}^n + r_{c2}^n, \quad (1)$$

where r_p is the radius of a parent artery, r_{c1} and r_{c2} are the radii of its two child arteries, and n is 2 for da Vinci's Rule and 3 for Murray's Law.

It is clear that characteristic arterial size is determined by dynamic remodelling of the vessel diameter in response to blood velocity near the wall (Langille, 1996; Simons et al., 2016). The force of blood flowing over the endothelial surface interacts with a junctional mechanosensory complex involving VEGFR protein that effects adjustments of wall dimensions to normalise the velocity near the wall (Baeyens et al., 2015). The normal 'set-point' is considered a regulated value of wall shear stress (WSS; dyn cm^{-2}). WSS can be evaluated from haemodynamic theory (Secomb, 2016). The famous haemodynamic paradigm that relates laminar flow rate to vessel size is the Poiseuille-Hagen (P-H)

Correspondence

Roger S. Seymour, School of Biological Sciences, Faculty of Sciences, University of Adelaide, Adelaide, 5005 SA, Australia.

E: roger.seymour@adelaide.edu.au

Accepted for publication 18 October 2019

Article published online 11 November 2019

equation, which is a solution to one of the incompressible Navier-Stokes equations:

$$\Delta P = \frac{(8 \dot{Q} \eta L)}{(\pi r_i^4)}, \quad (2)$$

where ΔP is the pressure difference required to achieve a rate of laminar flow (\dot{Q}) of a fluid of given viscosity (η) in a straight, horizontal tube of length (L) and inner radius (r_i). The pressure ($P = \text{force/area}$) driving flow acts on the cross-sectional area of the tube ($A = \pi r_i^2$), yielding a force (F_p):

$$F_p = A \Delta P = \frac{(\pi r_i^2)(8 \dot{Q} \eta L)}{(\pi r_i^4)} = \frac{(8 \dot{Q} \eta L)}{r_i^2}. \quad (3)$$

The flow creates a balancing frictional force (F_τ) which equals WSS (τ) occurring along the total inner surface area of the tube ($S = 2 \pi r_i L$):

$$F_\tau = S \tau = 2 \pi r_i L \tau. \quad (4)$$

because $F_p = F_\tau$, so

$$\tau = \frac{((8 \dot{Q} \eta L / r_i^2))}{2 \pi r_i L} = \frac{4 \dot{Q} \eta}{\pi r_i^3}. \quad (5)$$

Equation (5) is attributed to Poiseuille, but for distinction from the P-H equation, we have previously called it the 'shear stress equation' (Seymour et al., 2015, 2016). In imaging studies, WSS can also be calculated from the derivative of the velocity gradient near the wall, and the results largely concur with values calculated from the shear stress equation in the common carotid artery (Zarins et al., 1983). For physiological studies, Eq. (5) is more useful than the simplified version, $\tau = (r_i \Delta P) / (2 L)$, because ΔP and L are difficult to evaluate.

If, for any artery, $\dot{Q} \propto r_i^n$, then $\tau \propto r_i^{n-3}$. In particular, if $n = 3$, as in the case of Murray's Law, then $\dot{Q} \propto r_i^3$ and $\tau \propto r_i^0$; the exponent of 0 indicates that WSS is a constant throughout the arterial tree. Under da Vinci's Rule, $n = 2$ and so $\dot{Q} \propto r_i^2$ and $\tau \propto r_i^{-1}$; the negative exponent indicates that WSS decreases in larger vessels of the arterial tree.

The literature has conflicting information on the scaling of arterial size and blood flow rates. Much of the 20th century literature assumed that arterial WSS is constant at about 10–20 dyn cm⁻², implicitly affirming Murray's Law (Ku, 1997). However, natural selection does not necessarily conform to single factors, nor is it necessary for the value of the exponent n to be a constant throughout the arterial tree. In fact, n generally varies between 2 and 3 in experimental studies, but it is uncertain how n varies predictably in real arteries (Karau et al., 2001).

Murray's Law is upheld in studies of the small-medium arteries where the exponents n are nearly 3 for blood flow rate in relation to arterial diameter, and nearly zero for WSS. For example, flow rate varies with diameter to the power of 2.76 in human retinal arteries (Riva et al., 1985), similar to retinal arteries of rhesus monkeys (Zamir & Medeiros, 1982). Blood flow in cremaster muscle arteries of rats scales with diameter to the power of 3.01 (Mayrovitz & Roy, 1983). Blood flow in pial arteries on the surface of the brain of cats scales with diameter to the power of 2.98 (Kobari et al., 1984).

Conversely, da Vinci's Rule has received partial support by several investigators who have shown that WSS is not constant in the major arteries of humans and other mammals. In human studies, WSS varies between 2.9 to 15 dyn cm⁻² in the aorta and its immediate branches, and WSS decreases in arteries of larger radius at a power of -0.5 (Cheng et al., 2007). Interspecific studies also indicate that WSS is not a constant in individually named arteries (aortas and carotid arteries) of different-sized mammals, but decreases allometrically with increasing body mass to the approximate power of -0.2 (Greve et al., 2006; Cheng et al., 2007; Weinberg & Ethier, 2007).

Part of the variability in arterial radius, blood flow rate, and WSS undoubtedly relates to the fact that metabolic rates of some tissues are extremely variable, in particular the locomotory muscles. Flow rate in the femoral artery can increase 10-fold between rest and activity (Jorfeldt & Wahren, 1971); the femoral artery therefore shows low WSS at rest but, during activity, values are comparable to those in the similarly sized common carotid artery (Kornet et al., 2000). In contrast, brain perfusion is well autoregulated, varying less than approximately 20% regardless of whether a person is resting, physically or mentally exercising, or sleeping (Sokoloff et al., 1955; Armstrong, 1983; Hellström et al., 1996; Ogoh & Ainslie, 2009; Sato & Sadamoto, 2010; Hiura et al., 2014). The diameters of the cerebral arteries also change very little between rest and activity (Hellström et al., 1996) or in response to physiological changes in blood pressure and blood gas levels (Payne, 2016). Therefore, a tighter pattern of arterial variables, such as size, flow rate and WSS, may emerge in the circulation to the brain.

Nonetheless, there is disagreement about the scaling pattern in the major arteries of the human cerebral circulation. For example, a study that measured the lumen radii of the parent and branch arteries of the internal carotid, anterior cerebral and middle cerebral arteries from angiographs of healthy individuals found that the branching exponent n was 2.9 ± 0.7 (\pm SD), thus implying that Murray's Law prevails and WSS is constant (Rossitti & Löfgren, 1993). Another study of blood flow rate and cross-sectional area (A) of the human internal carotid artery and vertebral artery also supported Murray's Law, with flow being proportional to $A^{1.84}$, which corresponds to a very high $n = 3.74$ (Cebal et al., 2008). In contrast, the branching exponent for lumen

diameter in these three major cerebral arteries has been measured as 2.06, and the flow rate exponent as 2.45, thereby indicating that WSS increases in the smaller arteries and supporting da Vinci's Rule more than Murray's Law (Chnafa et al., 2017).

The present investigation is a meta-analysis of 19 imaging studies of blood flow rate and size in seven major cephalic arteries of humans. It analyses the relationships between all of the major arteries supplying the brain, and it represents the largest sample of studies that provide data for both flow rates and artery sizes in the same publication. It was gathered to test the applicability of Murray's Law and da Vinci's Rule in a system of relatively constant blood flow and to establish an empirical dataset useful for estimation of blood flow rate from measurements of cephalic arterial size in humans.

Methods

The literature was searched for studies involving *in vivo* measurements of major cephalic arteries from humans, including the common carotid (CCA), internal carotid (ICA), vertebral (VA), basilar (BA), anterior cerebral (ACA), middle cerebral (MCA) and posterior cerebral arteries (PCA). Techniques included phase-contrast magnetic resonance imaging (PC-MRI), transcranial Doppler ultrasound (TCD), multi-gate spectral Doppler ultrasound (MSDU), colour duplex Doppler ultrasound (CDDU) and high resolution ultrasound (HRU). Very many papers were screened, but studies were accepted only if they reported paired values for both blood flow rate and arterial lumen size under normal arterial blood pressure, so that WSS could be calculated independently for individual publications. There was no *a priori* reason to dismiss certain studies, so all data meeting these criteria were accepted as reported. Some variation may have resulted from differences in technique or image resolution, but there is no significant difference if exclusively PC-MRI studies are used in the analysis as compared with ultrasound studies (see Results). No weighting of studies with different sample sizes was undertaken to avoid pseudo-replication involving techniques, laboratories, subject populations, etc. The object of the meta-study was to treat an individual study as a single datum for each reported artery.

Blood flow rates were taken as reported means from presumably healthy adult individuals (control or disease-free groups). In a few cases, flow rate was calculated from mean velocity multiplied by cross-sectional lumen area. Arterial lumen radius was calculated from either diameter or cross-sectional area of *in vivo* images. WSS was calculated from each pair of data using the shear stress equation (Eq. 5), assuming a constant blood viscosity of $0.04 \text{ dyn s cm}^{-2}$ (Schmid-Schönbein et al., 1969; Amin & Sirs, 1985). This equation ignores actual viscosity at the wall, which may be slightly lower than the blood in general, as well as velocity changes during the cardiac cycle, blunted velocity profiles that occur in major arteries, and effects due to curves and bifurcations (Reneman et al., 2009). However, flow in the measured arteries is essentially laminar (Winkel et al., 2015) and effectively Newtonian (McGah & Capobianchi, 2015). If a full parabolic profile is assumed, according to Poiseuille flow conditions, the WSS may be underestimated (Dammers et al., 2003; Reneman et al., 2009). Measured *in vivo* velocity profiles for the human carotid artery that account for these factors yield WSS values of $11\text{--}13 \text{ dyn cm}^{-2}$ (Reneman & Hoeks, 2008; Reneman et al.,

2009). We found that the mean from seven common carotid artery WSS values in the present study was within this range ($12.01 \text{ dyn cm}^{-2}$), providing confidence that our method is accurate and valid to calculate the scaling exponent of WSS on arterial radius.

Values are presented as means with 95% confidence intervals (CI), and are compared with ANOVA calculated with Microsoft EXCEL add-in StatistixL (www.statistixl.com). Arithmetic data were plotted on arithmetic axes and power equations fitted with graphing and statistical software (GRAPHPAD Software Inc., La Jolla, CA, USA).

Results

Paired data for blood flow rate (\dot{Q} , $\text{cm}^3 \text{ s}^{-1}$) and arterial lumen radius (r_i , cm) were obtained from 19 imaging studies involving major arteries of the human cephalic circulation of healthy adult individuals (raw data available in Supporting information). The means and 95% CI of \dot{Q} , r_i and calculated WSS (τ , dyn cm^{-2}) are given in Table 1. Data for exclusively PC-MRI studies are presented in Supporting Information Table S2, but they are not significantly different from the entire set. For example, \dot{Q} in the ICA averaged $3.95 \pm 0.55 \text{ cm}^3 \text{ s}^{-1}$ in seven studies involving PC-MRI and $4.18 \pm 0.35 \text{ cm}^3 \text{ s}^{-1}$ in six studies that used ultrasound (*t*-test: $P = 0.52$). Other arteries were represented in fewer studies, also with no significant differences.

The contributions of each cephalic artery to total brain blood flow are summarised in Table 2. The two ICAs supply 75% and the two VAs 25%, of total brain blood flow. Single ICA flow ($4.05 \text{ cm}^3 \text{ s}^{-1}$) is slightly greater than the sum of single ACA + MCA flows ($3.62 \text{ cm}^3 \text{ s}^{-1}$), and slightly less than the sum of single ACA + MCA + PCA flows ($4.61 \text{ cm}^3 \text{ s}^{-1}$). However, the lower 95% confidence limit of individual ICA flow is $3.72 \text{ cm}^3 \text{ s}^{-1}$, which is greater than ACA + MCA flows, but less than ACA + MCA + PCA flows (Table 1). This indicates that, on average, the ICA contributes blood entirely to the ACA and MCA, and also partly to the PCA through the posterior communicating artery (PCoA) of the circle of Willis (Fig. 1). The VA also contributes to the PCA, as well as to the arteries of the upper spinal cord, brainstem and cerebellum. Thus the ICAs account for 88% and the VAs for 12% of cerebral perfusion.

The mean ICA flow ($4.05 \text{ cm}^3 \text{ s}^{-1}$) is about one-half that of the CCA ($8.09 \text{ cm}^3 \text{ s}^{-1}$), and the difference ($4.04 \text{ cm}^3 \text{ s}^{-1}$) should equal flow to the external carotid artery (ECA), which is nearly equal to ICA flow. However, published measurements of ECA flow rate are about 51–65% of that in the ICA on the same side (Schöning et al., 1994; Scheel et al., 2000; Hoi et al., 2010). The discrepancy is possibly attributed to branching of the superior thyroid artery, the lingual artery and the ascending pharyngeal artery from the ECA, which occurs close to its junction with the ICA (Fig. 1). It is possible that published measurements of ECA flow were made above this branching.

Calculated WSS is lowest in the larger CCA and VA, and highest in the smaller MCA and ACA (Table 1). A full-

Table 1 Mean \pm 95% CI arterial lumen radii, blood flow rates and calculated wall shear stress (WSS) of single arteries of the human cephalic circulation based on data from 19 PC-MRI and ultrasound imaging studies. Data for exclusively PC-MRI studies are given in Supporting information Table S1.

Artery Single	Number of studies <i>n</i>	Radius (r_i , cm)		Flow rate (\dot{Q} , cm ³ s ⁻¹)		WSS (τ , dyn cm ⁻²)	
		Mean	95% CI	Mean	95% CI	Mean	95% CI
Common carotid (CCA)	7	0.327	0.016	8.05	1.00	12.0	2.2
Internal carotid (ICA)	13	0.239	0.015	4.05	0.33	16.1	3.0
Basilar (BA)	3	0.190	0.021	2.39	0.29	18.2	4.1
Vertebral (VA)	8	0.186	0.014	1.37	0.21	11.3	3.0
Middle cerebral (MCA)	7	0.159	0.021	2.41	0.42	36.2	15.7
Posterior cerebral (PCA)	3	0.150	0.019	0.99	0.14	16.2	7.0
Anterior cerebral (ACA)	5	0.129	0.027	1.21	0.10	35.5	18.5

Table Mean total blood flow rates of the paired cephalic arteries of humans (= twice the single flow rates). Also shown are the percentage flow rates relative to total brain blood flow rate.

Variable	cm ³ s ⁻¹	cm ³ min ⁻¹	% total
Total brain flow rate	10.8	650	100
ICA \times 2	8.1	486	75
VA \times 2	2.7	164	25
(ACA + MCA) \times 2	7.2	434	67
(ACA + MCA+PCA) \times 2	9.2	553	85
PCoA \times 2*	0.9	53	8

*Posterior communicating artery (PCoA) flow is calculated (ICA – ACA – MCA) and its direction is from ICA to PCA. See Table 1 for other artery abbreviations.

factorial ANOVA and Student-Newman-Keuls *post hoc* test show that WSS is significantly higher ($P < 0.05$) in the MCA and ACA than in the other arteries (except for the MCA and BA, which are close to significant; $P = 0.07$). WSS in the MCA and ACA are not significantly different from each other ($P = 0.92$). WSS in the PCA is not significantly different from that in other arteries; however, the sample size for this artery is low ($n = 3$ studies).

Mean values of flow rate (\dot{Q} , cm³ s⁻¹) and WSS (τ , dyn cm⁻²) are plotted in relation to lumen radius (r_i , cm) in Fig. 2. Allometric power equations, derived from seven data points, for the cephalic arteries of humans are: $\dot{Q} = 82.6 r_i^{2.16 \pm 1.13}$ (95% CI) ($R^2 = 0.83$) and $\tau = 3.44 r_i^{-1.02 \pm 1.29}$ ($R^2 = 0.45$). Alternatively, polynomial regression equations ($\log \dot{Q} = -0.20 \log r_i^2 + 1.91 \log r_i + 1.82$, and $\log \tau = -0.20 \log r_i^2 - 1.09 \log r_i + 0.53$), derived from a larger analysis of 20 named arteries in nine species of mammals (Seymour *et al.* 2019), describe the human data well. Indeed, the sums of squared residuals of the logged flow rate data are nearly identical for the two models (0.115 for the interspecific polynomial regression equation and 0.107 for the human power equation). Focusing on patterns within individual

arteries, there is no obvious relationship between \dot{Q} and r_i (Supporting Information Fig. S1A), but a clear inverse relationship appears between WSS and r_i (Fig. S1B).

Discussion

Patterns of cranial blood flow

The primary motivation for this meta-analysis was to establish standard values for blood flow rate and arterial size in the major cephalic arteries of humans. This analysis includes 19 imaging studies of seven main arteries, compared with a previous meta-analysis of six studies of the common carotid artery only (Cheng *et al.*, 2007). Only one of the 19 studies included all seven arteries (Zhao *et al.*, 2015). Thus the present analysis goes as far as possible to reduce potential bias in individual studies and produce the best summary so far. The results are useful as a baseline against which future investigations can be compared, for example, clinical human studies and those concerning human evolution.

The literature indicates that there is little net flow of blood from one side of the brain to the other when the circle of Willis is complete. Previous studies show that flow rates are nearly identical in the left and right ICAs, but not necessarily in the VAs (Enzmann *et al.*, 1994; Schöning *et al.*, 1994; Sato & Sadamoto, 2010; van Ooij *et al.*, 2013; Zhao *et al.*, 2015). This aligns well with measurements of the nearly equal sizes of unpressurised cerebral arteries in the two hemispheres (Burlakoti *et al.*, 2017, 2018). Importantly, the ICA supplies the ACA and the MCA that service most of the cerebral cortex. The PCA receives blood mainly from the VA, supplemented by the ICA via the PCoA, and calculations based on our mean values indicate that the cerebrum receives 88% of its perfusion from the ICAs and only 12% from the VAs. This provides confidence that estimates of blood flow rates derived from ICA foramen size that capture the majority of cerebral perfusion and




Fig. 1 Major arteries of the healthy adult human cephalic circulation joining the Circle of Willis (cerebral arterial circle). Vessel diameters are drawn to scale for the major arteries examined in this study (dark red). Selected smaller arteries (pink; not to scale) are shown but were not directly measured, and other small cephalic arteries are omitted. Mean values of blood flow rate (\dot{Q} , $\text{cm}^3 \text{s}^{-1}$) are indicated next to each artery on one side only. Data are from Table 1.




Fig. 2 (A) Blood flow rate (\dot{Q} , $\text{cm}^3 \text{s}^{-1}$) and (B) calculated wall shear stress (τ , dyn cm^{-2}) in seven major cephalic arteries in relation to lumen radius (r_i , cm) in humans. Means and 95% confidence intervals are presented for the arteries, common carotid (CCA), internal carotid (ICA), basilar (BA), vertebral (VA), middle cerebral (MCA), posterior cerebral (PCA) and anterior cerebral (ACA). The solid lines are least square regressions from means of the present data, where $\dot{Q} = 82.6 r_i^{2.16}$ and $\tau = 3.44 r_i^{-1.02}$. The dotted lines represent polynomial regressions from previous analyses for arteries of mammals in general, where $\log \dot{Q} = -0.20 \log r_i^2 + 1.91 \log r_i + 1.82$, and $\log \tau = -0.20 \log r_i^2 - 1.09 \log r_i + 0.53$ (Seymour et al. 2019).

represent well cerebral metabolic rate in hominins among the primates (Seymour et al., 2015, 2016).

The low contribution by the VAs in humans is associated with variability in flow patterns among individuals. Flow rates in the PCoAs that connect the ICAs to the PCAs are small and apparently not essential. Flow along the PCoAs can be in either direction and is unpredictable in individuals that have either one or two of these arteries (Klötzsch et al., 1996; van Ooij et al., 2013). One or both of the PCoAs are absent in about 25% of humans (Lippert & Pabst, 1985). In contrast, absence of the ICAs is very rare, loss of one ICA occurring in less than 0.01%, and two ICAs in less than 0.001% of the population (Alexandre et al., 2016).

da Vinci’s Rule and Murray’s Law

The paradigms describing the circulation assume that it is a series of fractally bifurcating tubes. Although the major cephalic arteries do not conform to this model, but are rather a non-fractal, anastomosing network in which the arteries communicate in the circle of Willis (Fig. 1), we

assessed them as if they conformed, being partly justified by the fact that there is little net flow between hemispheres and cephalic flow rates are remarkably constant. Therefore, the working hypotheses are that the data for artery sizes and flow rates should have exponents of $n = 2$ for da Vinci’s Rule or $n = 3$ for Murray’s Law.

The exponent for blood flow rate in relation to lumen radius derived from the human cephalic arteries is 2.16 ± 1.13 (Fig. 2A), and the exponent for calculated WSS is -1.02 ± 1.29 (Fig. 2B). Although the 95% CI values include 3.0 or 0.0, respectively, the exponents are in much closer agreement to da Vinci’s Rule than to Murray’s Law. Increasing WSS in smaller cephalic arteries is similar to the results of a previous study of the ICA, MCA and ACA only (Chnafa et al., 2017). The two studies are also similar in showing a steep inverse relationship between WSS and radius between artery groups (Fig. S1B). We suspect that this is an artefact related to study-specific differences in measurement of lumen radius. Radii of all of these arteries are always highest in two studies (Enzmann et al., 1994; Zhao et al., 2015) and always lowest in three studies (Fahrig

et al., 1999; Chnafa et al., 2017; Bouillot et al., 2018), yet the flow rates are similar. This underscores the value of this meta-study. There are investigations of size and flow in human cephalic arteries that show similar increases in WSS in smaller vessels but do not comment specifically on da Vinci's Rule or Murray's Law (Enzmann et al., 1994; Fahrig et al., 1999; Reymond et al., 2012; van Ooij et al., 2013; Zhao et al., 2015). All of these studies are contrary to the conclusions of constant WSS in these vessels (Rossitti & Löfgren, 1993) and the cubic scaling of flow and diameter of the VA and ICA (Cebal et al., 2008).

The power equations set to the data in this analysis conform well to recent mathematical theory and to empirical results from mammals in general. In particular, the exponent of 2.16 for flow rate on radius (Fig. 2A) is similar to the exponent of 2.33 expected by recent haemodynamic theory of fractal-branching arterial trees (Huo & Kassab, 2012; Huo & Kassab, 2016). The present data are a subset of a larger study of arterial flow rate and WSS on lumen radius derived from 20 named systemic arteries among nine species of mammals (Seymour et al. 2019). That study confirms that a single exponent should not be applied to a broad range of arterial size and that a second-order polynomial equation best describes the log-transformed interspecific data. The derivative of the interspecific equation (the slope of the curve at any point) is equivalent to the exponent of a power equation at that point. The derivative is approximately 2.1 for the largest artery in this study (CCA $r_i = 0.327$ cm) and 2.3 for the smallest (PCA $r_i = 0.150$), so 2.16 is a reasonable exponent for these arteries. By comparison, the derivative is near 3.0 in very small arteries such as in the cremaster muscle of a rat, decreasing to about 2.0 in the aorta of a human (Seymour et al. 2019). Thus da Vinci's Rule may be applied to large arteries and Murray's Law to small ones.

The failure of the major arteries of the cephalic circulation to conform to Murray's Law does not entirely invalidate the idea that the structure of the circulatory system results in the least energy required by the heart. Our results are related to different functions of major and peripheral arteries. The major arteries, including those assessed in the present meta-analysis, buffer pulsatile pressures from the heart and minimise energy loss to reflected waves (Savage et al., 2008; Newberry et al., 2015), whereas the peripheral arteries distribute the blood according to the demands of the tissues and reduce the velocity to allow time for gas exchange (Zamir, 2001). It is clear that Murray's Law applies well in the peripheral arteries that are the major source of resistance to flow (Womersley, 1955; Sherman, 1981; Zamir et al., 1992; Savage et al., 2008; Reneman et al., 2009; McGah & Capobianchi, 2015). Low resistance in the major arteries is indicated by a small drop in mean blood pressure of approximately 5 mmHg between the aortic root and the MCA (Reymond et al., 2012). Murray

considered two factors to be important in affecting the structure of the circulatory system – energy costs associated with pumping blood and energy costs of building and maintaining the blood and vessels (Zamir, 1977). It is not clear which factor has more influence, with some modelling favouring pumping costs (Price et al., 2007) and others favouring material costs (Tekin et al., 2016). It is possible that the most important factor depends on the level of branching.

Conclusion

In summary, blood flow to the human brain is relatively constant and occurs through arteries that are usually bilaterally symmetrical. The ICAs supply 88% of the perfusion of the cerebral (cognitive) part of the brain. Among all of the seven major cephalic arteries, the relationship between blood flow rate and arterial size lies closer to da Vinci's Rule than to Murray's Law. Wall shear stress is not constant, but increases in smaller arteries supplying the brain.

Acknowledgements

We thank Professor Timothy J. Pedley, Associate Professor Jens Christian Brings Jacobsen and an anonymous referee for advice on the manuscript and Thomas Nelson for help with the literature. This study was supported by the Australian Research Council Discovery Project Grant DP170104952 to R.S.S.). Q.H. holds an Australian Postgraduate Award.

Author contributions

R.S.S. developed the original concept, gathered the literature and wrote the first draft. Q.H. and E.P.S. contributed to library research, data analysis and manuscript writing. All authors gave final approval of the manuscript.

Data Availability

The data are available in the supplementary material.

Electronic supplementary material

The raw data and their literature sources. Individual graphed data (Fig. S1). Results of PC-MRI data only (Table S1).

References

- Alexandre AM, Visconti E, Schiarelli C, et al. (2016) Bilateral internal carotid artery segmental agenesis: Embryology, common collateral pathways, clinical presentation, and clinical importance of a rare condition. *World Neurosurg* **95**, 620.e9–620.e15.
- Amin TM, Sirs JA (1985) The blood rheology of man and various animal species. *Q J Exp Physiol Cogn Med Sci* **70**, 37–49.

- Armstrong E** (1983) Relative brain size and metabolism in mammals. *Science* **220**, 1302–1304.
- Baeyens N, Nicoli S, Coon BG, et al.** (2015) Vascular remodeling is governed by a VEGFR3-dependent fluid shear stress set point. *Elife* **4**, e04645.
- Bouillot P, Delattre BMA, Brina O, et al.** (2018) 3D phase contrast MRI: Partial volume correction for robust blood flow quantification in small intracranial vessels. *Magn Reson Med* **79**, 129–140.
- Burlakoti A, Kumaratilake J, Taylor J, et al.** (2017) The cerebral basal arterial network: morphometry of inflow and outflow components. *J Anat* **230**, 833–841.
- Burlakoti A, Kumaratilake J, Taylor J, et al.** (2018) Asymmetries of total arterial supply of cerebral hemispheres do not exist. *Heliyon* **4**, e01086.
- Cebal JR, Castro MA, Putman CM, et al.** (2008) Flow-area relationship in internal carotid and vertebral arteries. *Physiol Meas* **29**, 585–594.
- Cheng C, Helderan F, Tempel D, et al.** (2007) Large variations in absolute wall shear stress levels within one species and between species. *Atherosclerosis* **195**, 225–235.
- Chnafa C, Bouillot P, Brina O, et al.** (2017) Vessel calibre and flow splitting relationships at the internal carotid artery terminal bifurcation. *Physiol Meas* **38**, 2044–2057.
- Dammers R, Stiff F, Tordoir JHM, et al.** (2003) Shear stress depends on vascular territory: comparison between common carotid and brachial artery. *J Appl Physiol* **94**, 485–489.
- Enzmann DR, Ross MR, Marks MP, et al.** (1994) Blood flow in major cerebral arteries measured by phase-contrast cine MR. *Am J Neuroradiol* **15**, 123–129.
- Fahrig R, Nikolov H, Fox AJ, et al.** (1999) A three-dimensional cerebrovascular flow phantom. *Med Phys* **26**, 1589–1599.
- Greve JM, Les AS, Tang BT, et al.** (2006) Allometric scaling of wall shear stress from mice to humans: quantification using cine phase-contrast MRI and computational fluid dynamics. *Am J Physiol Heart Circ Physiol* **291**, H1700–H1708.
- Hellström G, Fischer-Colbrie W, Wahlgren NG, et al.** (1996) Carotid artery blood flow and middle cerebral artery blood flow velocity during physical exercise. *J Appl Physiol* **81**, 413–418.
- Hiura M, Nariai T, Ishii K, et al.** (2014) Changes in cerebral blood flow during steady-state cycling exercise: a study using oxygen-15-labeled water with PET. *J Cereb Blood Flow Metab* **34**, 389–396.
- Hoi Y, Wasserman BA, Xie YYJ, et al.** (2010) Characterization of volumetric flow rate waveforms at the carotid bifurcations of older adults. *Physiol Meas* **31**, 291–302.
- Huo Y, Kassab GS** (2012) Intraspecific scaling laws of vascular trees. *J R Soc Interface* **9**, 190–200.
- Huo YL, Kassab GS** (2016) Scaling laws of coronary circulation in health and disease. *J Biomech* **49**, 2531–2539.
- Jorfeldt L, Wahren J** (1971) Leg blood flow during exercise in man. *Clin Sci* **41**, 459–473.
- Karau KL, Krenz GS, Dawson CA** (2001) Branching exponent heterogeneity and wall shear stress distribution in vascular trees. *Am J Physiol Heart Circulatory Physiol* **280**, H1256–H1263.
- Klötzsch C, Popescu O, Berlit P** (1996) Assessment of the posterior communicating artery by transcranial color-coded duplex sonography. *Stroke* **27**, 486–489.
- Kobari M, Gotoh F, Fukuuchi Y, et al.** (1984) Blood flow velocity in the pial arteries of cats, with particular reference to the vessel diameter. *J Cereb Blood Flow Metab* **4**, 110–114.
- Kornet L, Hoeks APG, Lambregts J, et al.** (2000) Mean wall shear stress in the femoral arterial bifurcation is low and independent of age at rest. *J Vasc Res* **37**, 112–122.
- Ku DN** (1997) Blood flow in arteries. *Annu Rev Fluid Mech* **29**, 399–434.
- Langille BL** (1996) Arterial remodeling: relation to hemodynamics. *Can J Physiol Pharmacol* **74**, 834–841.
- Lippert H, Pabst R** (1985) *Arterial variations in Man: Classification and frequency*. Munich: J.F. Bergmann.
- Mayrovitz HN, Roy J** (1983) Microvascular blood-flow: Evidence indicating a cubic dependence on arteriolar diameter. *Am J Physiol Heart Circulatory Physiol* **245**, 1031–1038.
- McGah PM, Capobianchi M** (2015) A modification of Murray's Law for shear-thinning rheology. *J Biomech Eng* **137**, e054503.
- Murray CD** (1926) The physiological principle of minimum work. I. The vascular system and the cost of blood volume. *Proc Natl Acad Sci USA* **12**, 207–214.
- Newberry MG, Ennis DB, Savage VM** (2015) Testing foundations of biological scaling theory using automated measurements of vascular networks. *PLoS Comput Biol* **11**, e1004455.
- Ogoh S, Ainslie PN** (2009) Cerebral blood flow during exercise: mechanisms of regulation. *J Appl Physiol* **107**, 1370–1380.
- van Ooij P van, Zwanenburg JJM, Visser F, et al.** (2013) Quantification and visualization of flow in the Circle of Willis: Time-resolved three-dimensional phase contrast MRI at 7 T compared with 3 T. *Magn Reson Med* **69**, 868–876.
- Payne S** (2016) *Cerebral Autoregulation: Control of Blood Flow in the Brain*. Basel: Springer Nature.
- Price CA, Enquist BJ, Savage VM** (2007) A general model for allometric covariation in botanical form and function. *Proc Natl Acad Sci USA* **104**, 13204–13209.
- Reneman RS, Hoeks APG** (2008) Wall shear stress as measured in vivo: consequences for the design of the arterial system. *Med Biol Eng Compu* **46**, 499–507.
- Reneman RS, Vink H, Hoeks APG** (2009) Wall shear stress revisited. *Artery Res* **3**, 73–78.
- Reymond P, Perren F, Lazeyras F, et al.** (2012) Patient-specific mean pressure drop in the systemic arterial tree, a comparison between 1-D and 3-D models. *J Biomech* **45**, 2499–2505.
- Richter JP** (1970) *The Notebooks of Leonardo da Vinci*. New York: Dover Publications.
- Riva CE, Grunwald JE, Sinclair SH, et al.** (1985) Blood velocity and volumetric flow rate in human retinal vessels. *Invest Ophthalmol Vis Sci* **26**, 1124–1132.
- Rossitti S, Löfgren J** (1993) Vascular dimensions of the cerebral arteries follow the principle of minimum work. *Stroke* **24**, 371–377.
- Sato K, Sadamoto T** (2010) Different blood flow responses to dynamic exercise between internal carotid and vertebral arteries in women. *J Appl Physiol* **109**, 864–869.
- Savage VM, Deeds EJ, Fontana W** (2008) Sizing up allometric scaling theory. *PLoS Comput Biol* **4**, e1000171.
- Scheel P, Ruge C, Schöning M** (2000) Flow velocity and flow volume measurements in the extracranial carotid and vertebral arteries in healthy adults: reference data and the effects of age. *Ultrasound Med Biol* **26**, 1261–1266.
- Schmid-Schönbein H, Wells R, Goldstone J** (1969) Influence of deformability of human red cells upon blood viscosity. *Circ Res* **25**, 131–143.
- Schöning M, Walter J, Scheel P** (1994) Estimation of cerebral blood flow through color duplex sonography of the carotid and vertebral arteries in healthy adults. *Stroke* **25**, 17–22.

- Secomb TW** (2016) Hemodynamics. *Comprehensive Physiology* **6**, 975–1003.
- Seymour RS, Angove SE, Snelling EP, et al.** (2015) Scaling of cerebral blood perfusion in primates and marsupials. *J Exp Biol* **218**, 2631–2640.
- Seymour RS, Bosiocic V, Snelling EP** (2016) Fossil skulls reveal that blood flow rate to the brain increased faster than brain volume during human evolution. *R Soc Open Sci* **3**, e160305.
- Seymour RS, Hu Q, Snelling EP, et al.** (2019) Interspecific scaling of blood flow rates and arterial sizes in mammals. *J Exp Biol* **222**, 199554.
- Sherman TF** (1981) On connecting large vessels to small. The meaning of Murray's Law. *J Gen Physiol* **78**, 431–453.
- Simons M, Gordon E, Claesson-Welsh L** (2016) Mechanisms and regulation of endothelial VEGF receptor signalling. *Nat Rev Mol Cell Biol* **17**, 611–625.
- Sokoloff L, Mangold R, Wechsler RL, et al.** (1955) Effect of mental arithmetic on cerebral circulation and metabolism. *J Clin Invest* **34**, 1101–1108.
- Tekin E, Hunt D, Newberry MG, et al.** (2016) Do vascular networks branch optimally or randomly across spatial scales? *PLoS Comput Biol* **12**, e1005223.
- Weinberg PD, Ethier CR** (2007) Twenty-fold difference in hemodynamic wall shear stress between murine and human aortas. *J Biomech* **40**, 1594–1598.
- Willis R** (2016) *On the Motion of the Heart and Blood in Animals*. Eugene: Resource Publications.
- Winkel LC, Hoogendoorn A, Xing RY, et al.** (2015) Animal models of surgically manipulated flow velocities to study shear stress-induced atherosclerosis. *Atherosclerosis* **241**, 100–110.
- Womersley JR** (1955) Method for the calculation of velocity, rate of flow and viscous drag in arteries when the pressure gradient is known. *J Physiol Lond* **127**, 553–563.
- Zamir M** (1977) Shear forces and blood vessel radii in the cardiovascular system. *J Gen Physiol* **69**, 449–461.
- Zamir M** (2001) Fractal dimensions and multifractality in vascular branching. *J Theor Biol* **212**, 183–190.
- Zamir M, Medeiros JA** (1982) Arterial branching in man and monkey. *J Gen Physiol* **79**, 353–360.
- Zamir M, Sinclair P, Wonnacott TH** (1992) Relation between diameter and flow in major branches of the arch of the aorta. *J Biomech* **25**, 1303–1310.
- Zarins CK, Giddens DP, Bharadvaj BK, et al.** (1983) Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res* **53**, 502–514.
- Zhao XX, Zhao MD, Amin-Hanjani S, et al.** (2015) Wall shear stress in major cerebral arteries as a function of age and gender – a study of 301 healthy volunteers. *J Neuroimaging* **25**, 403–407.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Fig. S1. (A) Blood flow rate (\dot{Q} , cm³ s⁻¹) and (B) calculated wall shear stress (τ , dyne cm⁻²) in seven major cephalic arteries in relation to lumen radius (r_i , cm) in humans. The arteries are the common carotid (CCA), internal carotid (ICA), vertebral (VA), basilar (BA), anterior cerebral (ACA), middle cerebral (MCA) and posterior cerebral (PCA).

Table S1. Data for cephalic artery size, blood flow rate and wall shear stress taken from 19 published studies of adult humans.

Table S2. Mean \pm 95% CI arterial lumen radii, blood flow rates, and calculated wall shear stress (WSS) of single arteries of the human cephalic circulation based on data from 10 exclusively PC-MRI studies.