




Dimethylformamide is a novel nitrilase inducer in *Rhodococcus rhodochrous*

V. P. Chhiba-Govindjee^{1,2} · K. Mathiba¹ · C. W. van der Westhuyzen¹ · P. Steenkamp¹ · J. K. Rashamuse³ · S. Stoychev¹ · M. L. Bode² · D. Brady² 

Received: 21 April 2018 / Revised: 20 August 2018 / Accepted: 3 September 2018 / Published online: 22 September 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Nitrilases are of commercial interest in the selective synthesis of carboxylic acids from nitriles. Nitrilase induction was achieved here in three bacterial strains through the incorporation of a previously unrecognised and inexpensive nitrilase inducer, dimethylformamide (DMF), during cultivation of two *Rhodococcus rhodochrous* strains (ATCC BAA-870 and PPPPB BD-1780), as well as a closely related organism (*Pimelobacter simplex* PPPPB BD-1781). Benzonitrile, a known nitrilase inducer, was ineffective in these strains. Biocatalytic product profiling, enzyme inhibition studies and protein sequencing were performed to distinguish the nitrilase activity from that of sequential nitrile hydratase-amidase activity. The expressed enzyme, a 40-kDa protein with high sequence similarity to nitrilase protein Uniprot Q-03217, hydrolyzed 3-cyanopyridine to produce nicotinic acid exclusively in strains BD-1780 and BD-1781. These strains were capable of synthesising both the vitamin nicotinic acid as well as β -amino acids, a compound class of pharmaceutical interest. The induced nitrilase demonstrated high enantioselectivity (> 99%) in the hydrolysis of 3-amino-3-phenylpropanenitrile to the corresponding carboxylic acid.

Keywords *Rhodococcus rhodochrous* · *Pimelobacter simplex* · Nitrilase · Induction · Expression · Dimethylformamide · Biocatalysis

Introduction

Nitrilase enzymes (EC 3.5.5.1) belong to a superfamily of enzymes that are found widely distributed in nature (Brenner 2002), and consist of proteins constituted of homo-dimeric building blocks with $\alpha\beta\beta\alpha$ - $\alpha\beta\beta\alpha$ sandwich folds that contain an active site displaying conserved cysteine, glutamic acid and lysine residues. C-site interactions between nitrilase

homodimers generate an unusual spiral polymer structure that may only then be catalytically active (Thuku et al. 2009). Nitrilases catalyse the hydrolysis of aliphatic, aromatic as well as aryl-aliphatic nitriles, generating a carboxylic acid product with the concomitant release of ammonia (Fig. 1), and are therefore of commercial value as biocatalysts in synthetic chemistry (Gong et al. 2012; Martínková et al. 2017).

A parallel nitrile hydrolysing reaction involving sequential action of a nitrile hydratase and amidase may be present in the same organism (Fig. 1). Unlike nitrile hydratase (EC 4.2.1.84), which is often constitutively expressed (Cramp and Cowan 1999; Nagasawa et al. 1986; Yamada and Kobayashi 1996; Endo and Watanabe 1989) (Table S1), nitrilase usually requires induction (Robinson and Hook 1964; Harper 1977; Lévi-Schil et al. 1995) with only a handful of strains demonstrating constitutive nitrilase enzyme production (Zheng et al. 2008, 2011; Gradley et al. 1994; Stalker et al. 1988; Gong et al. 2012; Fang et al. 2015; Ahmed et al. 2011) (Table S2). A number of nitrile-containing nitrilase inducers have been discovered, including benzonitrile, which was found to be an effective nitrilase enzyme inducer in *Alcaligenes faecalis* MTCC 10757 (Nageshwar et al. 2011);

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00253-018-9367-9>) contains supplementary material, which is available to authorized users.

✉ D. Brady
dean.brady@wits.ac.za

V. P. Chhiba-Govindjee
vchhiba@csir.co.za

¹ CSIR Biosciences, PO Box 395, Pretoria 0001, South Africa

² Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, Johannesburg PO Wits 2050, South Africa

³ The Department of Science and Technology (DST), Meiring Naude Road, Brummeria, Pretoria 0001, South Africa




Fig. 1 Schematic diagram showing nitrilase activity in comparison to the nitrile hydratase-amidase coupled reaction pathway

acetonitrile in *Stenotrophomonas maltophilia* AC21 (Badoei-Dalfard et al. 2016); 2-cyanopyridine in *Fusarium solani* IMI96840 (Vejvoda et al. 2010); tetrachloroterephthalonitrile (He et al. 2014) and propionitrile (Prasad et al. 2007) in *Rhodococci*; as well as isovaleronitrile in *Rhodococcus rhodochrous* J1 (Kobayashi et al. 1989) and *Nocardia guberula* NHB-2 (Sharma et al. 2011) (Table S2). However, these nitrile-based inducers may be subject to hydrolysis by the enzyme they induce, thereby inactivating them, and hence other classes of inducers are of interest. ϵ -Caprolactam, although ineffectual in organisms such as *Nocardia guberula* NHB-2 (Sharma et al. 2011), has been shown to induce nitrilase expression in *R. rhodochrous* J1 (Nagasawa et al. 1990) and *Fusarium proliferatum* (Yusuf et al. 2013) AUF-2.

Biocatalytic studies of nitrile hydrolysis in wild-type organisms are complicated by a number of factors that hinder unambiguous identification of the responsible enzyme system. Firstly, there are the two competing biocatalytic systems (nitrilase and nitrile hydratase-amidase) that can generate the same carboxylic acid product (Fig. 1). In many microorganisms, including *Rhodococci*, both of these enzyme systems may be located together on the genome (Martínková et al. 2009) and may also be constitutively expressed. Secondly, a nitrilase may be expressed but remain inactive if not oligomerised (Thuku et al. 2009). Thirdly, the enzymes are promiscuous: the formation of the amide intermediate, typically resulting from the action of a nitrile hydratase, may be the product of a nitrilase under certain conditions where the double hydrolysis reaction does not go to completion (Fernandes et al. 2006; Mateo et al. 2006); while a highly purified *R. rhodochrous* J1 amidase has been shown to slowly hydrolyse benzonitrile to benzoic acid (Kobayashi et al. 1998). Hence, a combination of chemical and molecular analysis is required to elucidate the responsible biocatalyst.

Herein, we demonstrate that dimethylformamide (DMF) is a cheap and effective nitrilase inducer in *Rhodococcus rhodochrous* and the closely related *Pimelobacter simplex*. Moreover, we definitively determined that nitrilase was responsible for nicotinic acid production using combined biocatalytic reaction profiling, inhibition studies and molecular

analysis. This is the first time that DMF has been shown to induce nitrilase.

Materials and methods

Isolates Three bacterial strains, namely *Rhodococcus rhodochrous* ATCC BAA-870, *R. rhodochrous* strain PPPPB BD-1780 as well as *Pimelobacter simplex* PPPPB BD-1781, were assessed in the current study. The two PPPPB strains are deposited at the Plant Pathogenic and Plant Protecting Bacteria Culture Collection (ARC, Pretoria, South Africa) while the ATCC strain is deposited at the American Type Culture Collection (Virginia, USA). BAA-870 was previously isolated from an industrial soil sample (Brady et al. 2004) and strains BD-1780 and BD-1781 were previously isolated from agricultural soil samples in the Gauteng province, South Africa (Rapheeha et al. 2017). All cells were cryopreserved in phosphate buffer (100 mM, pH 7.2) containing a 20% (v/v) final glycerol concentration.

Cell cultivation and induction All three strains were inoculated into tryptone soy broth (TSB) media for 72 h at 28 °C to allow for biomass production (0.5% inoculum volume, agitation at 180 rpm). Aseptic cell harvesting was done by centrifugation (10,000×g, 15 min) prior to biomass transfer into minimal media (Nagasawa et al. 1990) (control) or induction minimal media containing either 5 mM benzonitrile (BZN) or 0.5% v/v DMF (99%, Merck). All induction flasks were incubated for up to 72 h at 28 °C prior to cell harvesting by centrifugation (10,000×g, 15 min) and subsequent storage of the pellet at −20 °C.

Subsequent hydrolysis reactions were carried out in 1 mL with 100 mg wet weight cell paste (21 mg dry cell wt) in the presence of 4.16 mg (40 mmol/L) of 3-CP and agitation at 28 °C for 1 h.

Biocatalysis reactions 3-Cyanopyridine (3-CP) biocatalysis reactions involved reacting 200 mg wet weight cell paste (42 mg dry wt) in 4 mL of 0.1 M TRIS buffer, pH 9.0) cell preparation with 75.0 mg 3-CP in 1 mL methanol (reaction final volume 5 mL) with agitation at 28 °C for 15 min. Reaction quenching was by sample acidification with 1 M HCl.

Product recovery To extract nicotinamide, the reaction mixture was basified using a 1 M sodium hydroxide solution to a pH of 10.0 and then extracted (3× sample volume of ethyl acetate) in a mini-separation funnel. The combined organic phase was dried over anhydrous magnesium sulphate prior to product concentration by rotary evaporation. Flash column chromatographic purification allowed for the separation of the nitrile and the amide product.

Nicotinic acid extraction was done using a 1 M hydrochloric acid solution to acidify the reaction mixture to a final pH of 3.5–4 to protonate the carboxylic acid, followed by extraction into an organic phase (3× sample volume of *n*-butanol). Product concentration was achieved by removal of *n*-butanol in vacuo.

Sample preparation for NMR analysis Nicotinamide and nicotinic acid samples were solubilised in 200 µL of deuterated methanol prior to the addition of 400 µL of deuterated chloroform. Samples were analysed using a 400-MHz Varian Inova NMR spectrometer operating at 399.939 MHz (^1H) and 100.574 MHz (^{13}C) at 30 °C. Spectra were referenced against tetramethylsilane (TMS, δ 0.00 ppm) for ^1H NMR, and against residual protonated methanol (δ 49.1 ppm) for ^{13}C NMR, see Figs S1.1–8 for NMR spectra.

HPLC analysis method Nicotinic acid, nicotinamide and 3-CP present in the reaction mixtures were quantified by HPLC using a Waters Agilent 1100 (USA) system equipped with a binary pump, auto sampler and column thermostat. Data processing was done using Chemstation chromatography management data software. Separation was achieved using a C18 reverse phase column (Waters Spherisorb 5 cm ODS2 column (250 mm × 4.6 mm)), temperature controlled at 30 °C. The volume injected onto the column was 10 µL. The mobile phase was a binary gradient prepared from aqueous KH_2PO_4 solution (10 mM, pH 2.8) against acetonitrile solution (HPLC grade). An initial ratio of 90:10 (% v/v) was run for 6.0 min. Then, the gradient was ramped to a 30:70 (% v/v) aqueous:organic gradient for 5 min. The flow rate used was 1.0 mL/min (Binfeng et al. 2011). Detection was at 230 nm.

Calibration curves were constructed for nicotinic acid (1–20 mM), nicotinamide (1–10 mM) and 3-CP (1–5 mM) with correlation coefficients (R^2) of 0.990, 0.998 and 0.995, respectively. The retention times for nicotinic acid, nicotinamide and 3-CP were 3.541, 3.955 and 10.660 min, respectively.

Enantioselective biocatalytic resolution of β -aminonitriles

DMF-induced cells (100 mg wet wt) were re-suspended in a 0.1 M TRIS buffer solution (pH 9) prior to the addition of 10 mM of 3-amino-3-phenylpropanenitrile or 3-amino-3-(4-methoxyphenyl)propanenitrile. The reaction mixture (final volume 1 mL) was stirred at 28 °C with amide formation monitored by TLC. Biocatalytic reactions were terminated by acidification of the reaction (1 M HCl) prior to cell centrifugation (10,000×g, 5 min). The supernatant fraction was analysed by chiral HPLC.

Chiral HPLC was performed using a Hewlett Packard 1100 system consisting of a binary pump, autosampler, column thermostat compartment, diode array detector and Chemstation chromatograph data management software, version B.03.02. The analytical column used for separation was a

Chiralpak AD-H, 250 mm × 4.6 mm, 5 µm (Chiral Technologies Europe, France), conditioned for 1 h. The mobile phase used was a mixture of *n*-hexane and isopropanol (both analytical grade) in the volumetric ratio of 90:10. The flow rate used was 1.00 mL/min; column temperature was set at 25 °C. Detection was at 210 nm for 30 min.

Mass spectrometric analysis

Ultra-performance liquid chromatography mass spectrometry

A Waters UPLC coupled in tandem to a Waters SYNAPT G1 HDMS mass spectrometer was used to generate accurate mass data. Optimisation of the chromatographic separation was done utilising a Waters HSS T3 C18 column (150 mm × 2.1 mm, 1.7 µm) with the column temperature controlled at 60 °C. A binary solvent mixture was used consisting of aqueous formic acid (10 mM) (eluent A) (natural pH of 2.3) and 10 mM formic acid in methanol (eluent B). The initial conditions were 98% eluent A at a flow rate of 0.4 mL/min which was maintained for 5 min, followed by a linear gradient to 5% eluent A at 6 min. The conditions were kept constant for 1 min and then re-set to the initial conditions. The run time was 10 min and the sample injection volume was 1 µL.

Time-of-flight mass spectrometric analysis (ToF-MS)

A SYNAPT G1 mass spectrometer was used in V-optics and operated in electrospray mode. Leucine enkephalin (50 pg/mL) was used as reference calibrant to obtain typical mass accuracies between 1 and 5 mDa. The mass spectrometer was operated in ESI positive and negative mode with a capillary voltage of 4.0 kV, the sampling cone at 10 V and the extraction cone at 3.0 V. The scan time was 0.2 s covering the 50 to 650 Da mass ranges. The source temperature was 120 °C and the desolvation temperature was set at 450 °C. Nitrogen gas was used as the nebulisation gas at a flow rate of 550 L/h and cone gas was added at 50 L/h. MassLynx 4.1 (SCN 872) software was used to control the hyphenated system and for data manipulation.

Enzyme inhibition This was based on the inhibition of reactions converting 3-CP to nicotinic acid. Reactions were carried out using 100 mg of DMF-induced cells (wet wt) with nitrile degrading activity quantification performed using 20 mM 3-CP as substrate (final reaction volume 1 mL). All reactions were incubated at 28 °C for 60 min. Hydroxylamine (10 mM) (Prasad et al. 2009), ascorbic acid (10 mM) (Borsook et al. 1937) and nickel chloride (5 mM) were added to separate reactions to inhibit nitrile hydratase activity. Urea (5%) (Gregoriou and Brown 1979; Thalenfeld and Grossowicz 1976) and DEPA (7 mM) (Bauer et al. 1998; Brady et al. 2004) were included into reactions as aliphatic amidase inhibitors. Nitrilase competitive inhibitors benzylamine and benzaldehyde (10 mM) (Stevenson et al. 1992) were used to

confirm the presence of nitrilase activity. Reactant concentrations were determined by HPLC. Inhibition was determined as the percentage decrease in the yield of nicotinic acid compared to the control reaction.

Protein isolation and SDS-PAGE analysis

Cell disruption All control and induced cells were disrupted using a modified cell lysis method based on the protocol by Goncalves et al. (2006). Approximately 200 mg cell paste (wet wt) of the control, BZN and DMF-induced resting cells (72 h old induced cells) were re-suspended in 2 mL TRIS buffer (0.1 M, pH 8.8). Sodium dodecyl sulphate was added to the samples to a final concentration of 2.5% (v/v), prior to the addition of a protease inhibitor cocktail tablet into each sample [cOmplete ULTRA tablets, mini EDTA-free, Cat No.: 05892791001]. Glass beads (0.5 mm diameter, BioSpec Products, Cat No.: 11079105) were added into each tube to a final volume of 5 mL, with each subjected to 5 cycles of vortexing for 1 min followed by storage on ice for 1 min. Samples were subjected to vortexing and heating at 64 °C for 10 cycles of 1 min each. The cell lysate was centrifuged (13,000×g, 5 min) prior to storage of the samples at –20 °C. Further characterisation of the cell lysate was then done by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE).

Protein sample denaturation Protein samples were incubated at 70 °C for 10 min in the presence of a 1× reducing agent (Bolt™ sample reducing buffer, Cat No.: 1772696) and a 1× loading sample buffer (Bolt™ LDS sample loading buffer, Cat No.: 1772823) prior to protein separation on a commercially prepared denaturing SDS-PAGE gel.

SDS-PAGE gel profiling Denaturing SDS-PAGE gel electrophoresis was done using the method of Laemmli (1970). A BSA standard curve (100–800 ng/10 µL) was used for quantification purposes. All samples were loaded onto a precast 4–12% PAGE gel [Bolt™ 4–12% Bis-TRIS Plus, Cat No.: NW04125, LTC scientific] with each sample undergoing denaturation in the presence of a reducing agent at 70 °C for 10 min prior to loading onto the gel, and running under constant voltage conditions (200 V) using a commercial morpholinoethanesulfonic acid buffer [Bolt™ MES buffer, Cat No.: 1771547]. Electrophoresis was performed at room temperature until the loading dye had migrated to the bottom of the gel. Gel staining was done using a Coomassie B-250 stain solution. Molecular weight determination was done using the Fermentas Pst1 molecular weight marker (ThermoFischer, PageRuler Plus Prestained Protein ladder-Thermoscientific, Cat No.: 26634). The Biorad Chemi Doc™ imaging system with Image Lab 4.1 software was used to determine the intensity of the bands and quantified against a

BSA standard curve. Bands were excised from each of the protein gels and submitted for in-gel digestion prior to LC-MSMS sequence analysis.

Sample preparation for peptide sequencing by LC-MSMS

In-gel digestions SDS-PAGE excised gel slices were diced into ~1 mm² fragments, with 200 µL of 50 mM NH₄HCO₃ in 50% methanol being added into each sample, and vortexed for 20 min before discarding the supernatant. This was repeated to allow for adequate gel de-staining prior to the addition of 100 µL of 75% acetonitrile. The resulting sample was then vortexed for 20 min with the supernatant being discarded and the gel pieces dried under vacuum. Protein plugs were stored at –20 °C prior to sample denaturation and alkylation.

Alkylation Approximately 25 µL of freshly prepared 10 mM dithiothreitol (DTT) prepared in 25 mM NH₄HCO₃ was added to each gel slice prior to vortexing. The protein sample was then incubated at 60 °C for 1 h to allow for sample denaturation and reduction before being cooled to room temperature. Approximately 500 µL of MS grade acrylonitrile was added to the samples, which were incubated at room temperature for a further 10 min with the supernatant being discarded. Approximately 25 µL of a 55 mM iodoacetamide solution prepared in 25 mM NH₄HCO₃ was added to the protein gel pieces before vortexing and incubation at room temperature for 20 min. The supernatant was discarded and the gel pieces were washed with 100 µL NH₄HCO₃ prepared in milli-Q water. The supernatant was once again discarded and the individual gel pieces dehydrated by the addition of 100 µL of a 50% acrylonitrile in 25 mM NH₄HCO₃ solution. Protein plugs were dried under vacuum and stored at –20 °C.

Trypsin digestion This was performed by incubating the protein plug with 1–5 ng/µL MS grade trypsin. Approximately 50 µL of 25 mM NH₄HCO₃ was added to each sample prior to incubation overnight at 37 °C to facilitate protein cleavage. Peptide extraction was carried out in protein Lo-Bind tubes (Eppendorf, Cat No.: 0030108035). Approximately 30 µL of 50% acrylonitrile/5% formic acid in milli-Q deionised water was added to the protein plugs with the samples being vortexed for 30 min at room temperature. All samples were centrifuged and then sonicated for 5 min prior to collection of the digest solution and vacuum drying before analysing by LCMS-MS.

LCMS-MS analysis MS-derived proteomics analysis was done using the CompOmics PeptideShaker programme (Vaudel et al. 2015). Digests were re-suspended in 20 µL of 2% acetonitrile/0.2% formic acid and analysed using a Dionex Ultimate 3000 RSLC system coupled to a Sciex 6600 TripleTOF mass spectrometer. Peptides were first de-salted

on an Acclaim PepMap C18 trap column (100 $\mu\text{m} \times 2 \text{ cm}$) for 2 min at 15 $\mu\text{L}/\text{min}$ using 2% acetonitrile/0.2% formic acid, and then separated on an Acclaim Pepmap C18 RSLC column (300 $\mu\text{m} \times 15 \text{ cm}$, 2 μm particle size). Peptide elution was achieved using a flow rate of 8 $\mu\text{L}/\text{min}$ with a gradient of 4–60% B in 15 min (A: 0.1% formic acid; B: 80% acetonitrile/0.1% formic acid). An electrospray voltage of 5.5 kV was applied to the emitter. The 6600 TripleTOF mass spectrometer was operated in Data Dependant Acquisition mode. Precursor MS scans were acquired from m/z 400–1500 using an accumulation time of 250 ms followed by 30 MS-MS scans, acquired from m/z 100–1800 at 100 ms each, for a total scan of 3.3 s. Multiply charged ions (2+, 5+, 400–1500 m/z) were automatically fragmented in Q2 collision cells using nitrogen as the collision gas. Collision energies were chosen automatically as a function of m/z and charge.

Protein pilot v5.01.10 using Paragom search engine (Sciex) was used for comparison of the obtained MS/MS spectra against protein sequences in a Uniprot Swissprot 2016 database. Proteins with a threshold above $\geq 99.9\%$ confidence were reported. Peptides with $\geq 95\%$ confidence were used to report sequence coverage.

Results

Induction studies

Cultures of the strains *R. rhodochrous* ATCC BAA-870, *R. rhodochrous* BD-1780 and *P. simplex* BD-1781 were suspended in a minimal medium and induced with either benzonitrile (BZN) or dimethylformamide (DMF). After incubation for set periods, 100 mg wet wt cell paste of these cultures was used for a 1 h biocatalytic hydrolysis of 4.16 mg 3-cyanopyridine (3-CP; Fig. 2), and the products were identified and quantified by LC-MSMS and ultra-performance liquid chromatography mass spectrometry (UPLC-MS) in ESI⁺ mode (Fig. S2).

For all three strains, induction with DMF resulted in the production of nicotinic acid from 3-CP. In the absence of DMF, no nicotinic acid was formed whatsoever. Control cultures and BZN-induced cells of strains BD-1780 and BD-1781 generated only small amounts of nicotinamide, with large amounts of unreacted 3-CP remaining. However, for strain BAA-870, both control and BZN-induced cells showed complete conversion of 3-CP to nicotinamide (Fig. 3).

Biocatalytic reactions

To generate sufficient product for complementary analytical procedures (chromatography and NMR), larger reactions were performed (200 mg wet wt biocatalyst and 75 mg 3-CP in 5 mL for 15 min). These revealed that the biocatalytic reactions of




Fig. 2 3-Cyanopyridine (1), nicotinamide (2) and nicotinic acid (3), dimethylformamide (4), benzonitrile (5) and ϵ -caprolactam

DMF-induced strain BD-1780 and BD-1781 cells generated only nicotinic acid after complete conversion of 3-CP. However, the DMF-induced BAA-870 cells also generated nicotinamide in addition to nicotinic acid products in these more concentrated reactions, possibly due to greater substrate availability for the nitrile hydratase. The product identity was confirmed using ¹H and ¹³C NMR (Fig. S1; Table S3) as well as HPLC-MS. The isolated nitrile, amide and acid yields for each reaction were determined by weighing after chromatographic separation (Tables 1 and S4). The specific nitrilase activity of strains BD-1780 and BD-1781 on 3-cyanopyridine was calculated as 0.164 and 0.202 $\mu\text{mol}/\text{min}/\text{mg}$ DCW, respectively.

High enantiomeric excess of a product of biocatalytic hydrolysis of a racemate is indicative of a single enzyme system and thus enantioselectivity of the biocatalysts was investigated. Reaction of racemic 3-amino-3-phenylpropanenitrile and 3-amino-3-(4-methoxyphenyl)propanenitrile in the presence of DMF-induced cells of strains BD-1780 and BD-1781 resulted in enantioselective kinetic resolution of these substrates. Excellent enantiomeric excess of the residual nitrile was observed when cells of either strain were reacted with 3-amino-3-phenylpropanenitrile ($> 99\%$ ee), while moderate enantioselectivity was found in the case of 3-amino-3-(4-methoxyphenyl)propanenitrile (34–41%) (Table S5). Based on the previously determined formation of (*R*)-3-amino-3-phenylpropionic acid by these strains (Chhiba et al. 2014), the residual nitrile can tentatively be assigned as the (*S*)-enantiomer (Fig. 4). In comparison, the nitrile hydratase of BAA-870 enantioselectively hydrolysed the opposite (*S*)-enantiomer of 3-amino-3-(4-methoxyphenyl)propanenitrile to generate the (*S*)-amide, but was relatively non-enantioselective in the hydrolysis of 3-amino-3-phenylpropanenitrile (Chhiba et al. 2012).

Enzyme inhibition studies

The enzyme system responsible (nitrilase or nitrile hydratase-amidase) could not be definitively assigned solely by the




Fig. 3 Biocatalytic hydrolysis of 3-cyanopyridine. 3-Cyanopyridine (blue); nicotinamide (red); nicotinic acid (green). X-axis: induction period: 0 h (1); 24 h (2); 48 h (3) and 72 h (4). No analysis was made of BAA-870 cells at 24 h. All reactions were carried out for 1 h in 1 mL

with 100 mg dry wt cells and agitation at 28 °C in the presence of 40 mmol/L 3-CP. Average of six data points, consisting of triplicate flasks with duplicate samples per flask

reaction product profile as evidenced by the differences seen between small scale and large scale reactions. Consequently, reactions with DMF-induced cells were subjected to nitrilase, nitrile hydratase and amidase inhibitors (Table 2).

Nicotinic acid generation by strains BD-1780 and BD-1781 was considerably reduced in the presence of the nitrilase inhibitors benzaldehyde (83–90%) and benzylamine (69–82%) (Table 2), indicating that nicotinic acid production was dependent on nitrilase activity. Nicotinic acid production by BAA-870 was influenced to a lesser extent (9–22%), presumably due to the presence of an alternative route via nitrile hydratase (as demonstrated in the induction studies) and an amidase (detected in molecular studies below). Conversely, the nitrile hydratase and amidase inhibitors had a negligible effect, indicating that the nitrilase route was dominant, even in the case of BAA-870. Both benzaldehyde and benzylamine resemble nitrilase reaction intermediates, with benzylamine similar to the thio-imidate intermediate and benzaldehyde analogous to the thio-ester intermediate, and hence, a model for competitive inhibition can be proposed (Fig. 5).


Molecular characterisation

Inhibition studies undertaken on the DMF-induced cells suggested that the nitrile degrading activity of strains BD-1780 and BD-1781 was the result of an induced nitrilase enzyme. This, when coupled with the isolation of mixed amide and acid products from the DMF-induced cells of strain BAA-870, indicates that the nicotinic acid observed for BAA-870 could have been the result of an interplay of nitrilase, nitrile hydratase and amidase. Hence, it was decided to confirm the identity of the responsible enzymes through enzyme expression studies.

SDS-PAGE protein profiling was done on control, BZN and DMF-induced cells of all three strains (Fig. 6). With DMF induction, a dominant 40 kDa size protein, representing 11–12% of the total cytosolic soluble protein, was expressed in each case [BAA-870 (Fig. 6a lanes 11–12); strain BD-1780 (Fig. 6b lanes 6–7); strain BD-1781 (Fig. 6c lanes 6–7)]. No such band was observed in the control and BZN-induced cells. BAA-870 expressed an additional major protein at 25 kDa (Fig. 6a lanes 7–12).

Table 1 3-Cyanopyridine biocatalysis yields from DMF-induced cells

Strain	Isolated yield (%)			Implied enzyme system
	3-Cyanopyridine	Nicotinamide	Nicotinic acid	
BD-1780	0	0	80	Nitrilase
BD-1781	0	0	58	Nitrilase
BAA-870	0	22	38	Nitrile hydratase-amidase, nitrilase



R = H; 3-amino-3-phenylpropanenitrile

R = OCH₃; 3-amino-3-(4-methoxy-phenyl)propanenitrile

Fig. 4 Enzymatic kinetic resolution of racemic β -aminonitriles

Protein bands correlating to 100, 70, 55, 40 and 25 kDa were excised and analysed by LC-MSMS. The SDS-PAGE protein profiles for *P. simplex* strain BD-1781 cells showed smearing consistent with extensive proteolysis, with only faint bands discernable even with addition of a protease inhibitor cocktail during cell lysis, so bands were excised relative to the molecular weight markers. Reference to the MEROPS website showed that *P. simplex* cells produce an array of proteases, with around 82 known and putative proteolytic enzymes identified in this one strain (MEROPS the Peptidase Database – EMBL-EBI 2017), which may explain the general protein degradation.

Nitrilase peptide fragments corresponding to Uniprot Q-03217 were found in the 40 kDa protein band from DMF-induced cells of all three strains evaluated, with no nitrilase fragments being found in any of the control or BZN-induced cells. The protein fragments showed >99% sequence identity to an aliphatic nitrilase protein produced by *R. rhodochrous* J1 with 36.1% of protein coverage observed in BAA-870; 92.62% of the total sequence coverage located in the BD-1780; and 35.3% sequence coverage identified in the strain BD-1781 sample (Fig. S3).

The 25 kDa band in all the SDS-PAGE lanes of the cell lysates of BAA-870 cells was determined to be the alpha subunit of a constitutively expressed low molecular weight nitrile hydratase (Fig. S4). An amidase was also identified in




Fig. 5 Proposed competitive inhibition points for benzaldehyde and benzylamine in the nitrilase catalytic cycle based on structural similarity to the reaction intermediates

all cell lysates of BAA-870 (Fig. S5) as the 55 kDa band (P-84650, enantioselective amidase from *R. rhodochrous*, 6% sequence coverage, >99% sequence similarity) (Pertsovich et al. 2005). No nitrile hydratase or amidase protein sequence fragments were located in any of the strain BD-1780 and BD-1781 samples, suggesting the inducible nature of these enzymes in these strains. The characterisation of the three strains under DMF induction has been summarised in Table 3).

An array of transcription and translational proteins were found to be present in the lysates of the DMF-induced cells. These included both the 30S and 50S ribosomal proteins crucial for protein translation; DNA-directed RNA polymerase; various elongation factors (EF-U, EF-G as well as EF-Ts); as well as numerous 60 kDa chaperonin proteins. One chaperonin peptide identified in all three strains was Uniprot Q-2JA12 (Fig. S6), which showed high amino acid sequence similarity to Uniprot O50305 (GroEL 60 kDa chaperonin protein from *Bacillus halodurans*), which is associated with enhanced nitrilase activity (Petříčková et al. 2012).

Table 2 Inhibition of 3-CP to nicotinic acid by DMF-induced cells

Strain ^a	Nitrilase inhibition (%)		Nitrile hydratase inhibition (%)			Amidase inhibition (%)	
	Benzylamine	Benzaldehyde	Ascorbic acid	Hydroxylamine	NiCl ₂	Urea	DEPA
BD-1780	69	90	0	0	0	0	0
BD-1781	82	83	2	0	0	0	0
BAA-870	22	9	1	0	0	0	0

^a Reactions (1 h) were carried out using 20 mM 3-CP as substrate with 100 mg wet wt of resting cells. Results are expressed as the percentage reduction in the amount of nicotinic acid produced in the presence of the inhibitor




Fig. 6 SDS-PAGE gel of protein soluble fraction from induced and non-induced cells. BAA-870 (**a**): lane 1, PageRuler Plus Prestained Protein ladder; lanes 3–6, BSA standard concentrations 100–500 ng μ L protein; lanes 7–8, BAA-870 control cell sample; lanes 9–10, BAA-870 BZN-induced cell sample; lanes 11–12, BAA-870 DMF-induced cell sample.

Strain BD-1780 (**b**): lanes 2–3, BD-1780 control cell sample; lanes 4–5, BD-1780 BZN-induced cell sample; lanes 6–7, DMF-induced cell sample. Strain BD-1781 (**c**): lane 1, PageRuler; lanes 2–3, BD-1781 control cell sample; lanes 4–5, BD-1781 BZN-induced cell sample; lanes 6–7, DMF-induced cell sample

Discussion

DMF has not previously been assessed as a nitrilase inducer. Although organic solvents, including DMF, have previously been used to enhance nitrilase and nitrile hydratase activity in biocatalytic reactions (Stevenson et al. 1992; Nagasawa et al. 2000; Kaul and Banerjee 2008; Tauber et al. 2000), these effects were most likely related to solubility of the enzyme substrate during the reaction. In the present study, it was shown that *Rhodococcal* strains BD-1780 and BAA-870 and the closely related *P. simplex* strain BD-1781 expressed a DMF-inducible nitrilase.

DMF-induced cells of strains BD-1780 and BD-1781 were shown to convert 3-cyanopyridine (3-CP) solely to nicotinic acid. Addition of the nitrilase inhibitors benzaldehyde and benzylamine reduced 3-CP hydrolysis in agreement with Stevenson et al. (1992) who showed that these compounds competitively inhibited nitrilase activity in resting cells of *Rhodococcus* ATCC 39484. The results of biocatalysis and

enzyme inhibition studies were confirmed by SDS-PAGE and LC-MSMS amino acid sequence analysis which showed that DMF-induced cells expressed a dominant ~40 kDa protein with a sequence identical to a known nitrilase. BAA-870 showed additional constitutive expression of a nitrile hydratase that synthesised nicotinamide. Inhibitors for nitrile hydratase and amidase (Yamada and Kobayashi 1996; Endo and Watanabe 1989; Prasad et al. 2009; Nagasawa et al. 1991; Tani et al. 1989; Zangi et al. 2009; Precigou et al. 2004; Leonova et al. 2000; Bauer et al. 1998; Brady et al. 2004) had negligible effect, indicating that nitrilase was the primary nicotinic acid producer.

Nitrilase enzyme activation by subunit dimerization and subsequent polymerisation is often achieved through the action of specialised chaperone proteins. Nitrilase purification studies on *Bacillus pallidus* Dac521, *Comomonas testosteroni* and *Pseudomonas fluorescens* DSM 7155 showed that nitrilases are often co-purified with the chaperonin GroEL (Lévi-Schil et al. 1995; Almatawah et al. 1999; Layh et al.

Table 3 Summary of nitrile hydrolysing enzymes in DMF-induced cells

Strain	Product ^a	Inhibitors of nicotinic acid formation	Molecular characterisation ^b
BD-1780	Nicotinic acid	Nitrilase inhibitors (61.3–88.4%)	Nitrilase (Q03217) 40 kDa; 12% of total soluble protein
BD-1781	Nicotinic acid	Nitrilase inhibitors (79.5–87.5%)	Nitrilase (Q03217) 40 kDa; 11% of total soluble protein
BAA-870	Nicotinic acid and nicotinamide	Nitrile hydratase inhibitors (12–16%)	Nitrilase (Q03217) 40 kDa; 12% of total soluble protein; nitrile hydratase (P29378) LMW α -subunit 20 kDa; and amidase (P-84650) 55 kD)

^a Analysis by UPLC-MS; ¹H NMR; ¹³C NMR

^b SDS-PAGE and LC-MSMS

1998). Similar enhancement of activity was seen by co-expressing fungal nitrilases from *Gibberella moniliformis* and *Penicillium marneffei* with GroEL (Petříčková et al. 2012). In the present study, GroEL was co-expressed with nitrilase in strains BD-1780, BD-1781 and BAA-870.

The role that DMF plays in nitrilase induction in these cells is not obvious but, as DMF is toxic to most cells, nitrilase activity induction may be part of a cell detoxification mechanism. Nitrilase expression in plants is often to facilitate cellular detoxification when the breakdown of glucosinolates yields toxic isothiocyanates and nitrile compounds (Miller and Conn 1980). DMF (or a metabolite such as dimethylamine or *N*-methylformamide) could function as a *nitR* gene regulator. Nitriles, although the most common, are not the only inducers of nitrilase (Table S2). Herai et al. (2004) have demonstrated that a ϵ -caprolactam-induced *R. rhodochrous* J1 nitrilase expression system (Nagasawa et al. 1990) could be of value in industrial heterologous protein production. By superimposing ϵ -caprolactam and DMF, a structural similarity is evident (Fig. 7), implying that the mechanism of induction may be similar. However, DMF is commercially attractive at a fraction of the cost of ϵ -caprolactam. Preliminary results (SDS-PAGE) from our laboratory indicate that the structurally similar *N,N*-dimethylacetamide and *N*-methylpyrrolidone also induce nitrilase in strain BD-1780.

In conclusion, DMF is a novel inducer that can be used to selectively induce nitrilase but not nitrile hydratase. This permits the catalytic specificity inherent in the nitrilase to be expressed, unencumbered by the competing nitrile hydratase-amidase system. This is particularly desirable when resolving enantiomers or isomers as the competing nitrile biocatalytic systems may have opposing enantio- or regio-selectivity, potentially resulting in poor enantiomeric excess (ee) or isomer yields. Through application of multiple analytical




Fig. 7 Overlay after energy minimisation of DMF and ϵ -caprolactam (ChemDraw) showing the structural similarities of the two molecules

techniques, we were also able to unequivocally define which enzymes were responsible for biocatalytic nitrile hydrolysis and enantiomeric resolution in prior studies using these microbial strains (Chhiba et al. 2012, 2014; Brady et al. 2004; Rapheeha et al. 2017; Kinfel et al. 2009). These results expand the small group of non-nitrile nitrilase inducers previously only represented by the lactams reported by Nagasawa et al. (1990).

Acknowledgements We would like to thank the following people at the University of the Witwatersrand: Prof. Joseph Michael; Prof. Charles De Koning; and Dr. Amanda Rousseau. At the CSIR, we would like to extend thanks to Mr. Siphon Mamputhu (for assistance with LC-MSMS protein sequencing) as well as Dr. Priyen Pillay, Dr. Lusisiwe Kwezi, Dr. Ofense Poe, Advaita Singh, Sibongile Mtimka, Dr. Daria Rutkowska, Dr. Kevin Wellington and Dr. Tsepo Tsekoa.

Funding Support was received from the Department of Science and Technology *Biocatalysis Initiative* (Grant 0175/2013).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Ahmed K, Kumar M, MS KCG, Shaik TB (2011) Bioconversion of acrylonitrile to acrylic acid by *Rhodococcus ruber* strain AKSH-84. *J Microbiol Biotechnol* 21:37–42
- Almatawah QA, Cramp R, Cowan DA (1999) Characterization of an inducible nitrilase from a thermophilic bacillus. *Extremophiles* 3: 283–291
- Badoei-Dalfard A, Karami Z, Ramezani-pour N (2016) Bench scale production of nicotinic acid using a newly isolated *Stenotrophomonas maltophilia* AC21 producing highly-inducible and versatile nitrilase. *J Mol Catal B Enzym* 133:S552–S559
- Bauer R, Knackmuss H-J, Stolz A (1998) Enantioselective hydration of 2-arylpropionitriles by a nitrile hydratase from *Agrobacterium tumefaciens* strain d 3. *Appl Microbiol Biotechnol* 49:89–95
- Binfeng L, Jinhuan S, Tao J (2011) Enzyme and process development for production of nicotinamide. *Org Process Res Dev* 15:291–293
- Borsook H, Davenport HW, Jeffreys CEP, Warner RC (1937) The oxidation of ascorbic acid and its reduction in vitro and in vivo. *J Biol Chem* 117:237–279
- Brady D, Beeton A, Kgaje C, Zeevaart J, van Rantwijk F, Sheldon RA (2004) Characterisation of nitrilase and nitrile hydratase biocatalytic systems. *Appl Microbiol Biotechnol* 64:76–85
- Brenner C (2002) Catalysis in the nitrilase superfamily. *Curr Opin Struct Biol* 12:775–782
- Chhiba VP, Bode ML, Mathiba K, Kwezi W, Brady D (2012) Enantiomeric biocatalytic hydrolysis of β -aminonitriles to β -aminoamides using *Rhodococcus rhodochrous* ATCC BAA-870. *J Mol Catal B Enzym* 76:68–74
- Chhiba VP, Bode M, Mathiba K, Brady D (2014) Enzymatic stereoselective synthesis of β -amino acids. In: Riva S, Fessner WD (eds) *Cascade biocatalysis: integrating*

- stereoselective and environmentally friendly reactions. Wiley-VCH, Germany, pp 297–313
- Cramp RA, Cowan DA (1999) Molecular characterization of a novel thermophilic nitrile hydratase. *Biochem Biophys Acta* 1431:249–260
- Endo T, Watanabe I (1989) Nitrile hydratase of *Rhodococcus* sp. N774 purification and amino acid sequences. *FEBS Lett* 243:61–64
- Fang S, An X, Liu H, Cheng Y, Hou N, Feng L, Huang X, Li C (2015) Enzymatic degradation of aliphatic nitriles by *Rhodococcus rhodochrous* BX2, a versatile nitrile-degrading bacterium. *Bioresour Technol* 185:28–34
- Fernandes BCM, Mateo C, Kiziak C, Chmura A, Wacker J, van Rantwijk F, Stolz A, Sheldon RA (2006) Nitrile hydratase activity of a recombinant nitrilase. *Adv Synth Catal* 348:2597–2603
- Goncalves ER, Hara H, Miyazawa D, Davies JE, Eltis LP, Mohn WW (2006) Transcriptomic assessment of isozymes in the biphenyl pathway of *Rhodococcus* sp. strain RHA1. *Appl Environ Microbiol* 72:6183–6193
- Gong J-S, Lu Z-M, Li H, Shi JS, Zhou Z-M, Xu Z-H (2012) Nitrilases in nitrile biocatalysis: recent progress and forthcoming research. *Microb Cell Factories* 11:142
- Gradley ML, Deverson CJF, Knowles CJ (1994) Asymmetric hydrolysis of R(-)-S (+)-2-methylbutyronitrile by *Rhodococcus rhodochrous* NCIMB 11216. *Arch Microbiol* 161:246–251
- Gregoriou M, Brown PR (1979) Inhibition of the aliphatic amidase from *Pseudomonas aeruginosa* by urea and related compounds. *Eur J Biochem* 96:101–108
- Harper DB (1977) Microbial metabolism of aromatic nitriles: enzymology of aromatic nitriles: enzymology of C-N cleavage by *Nocardia* sp. (*Rhodococcus* group) N. C. I. B. 11216. *Biochem J* 165:309–319
- He Y-C, Wu Y-D, Pan X-H, Ma C-L (2014) Biosynthesis of terephthalic acid, isophthalic acid and their derivatives from the corresponding dinitriles by tetrachloroterephthalonitrile-induced *Rhodococcus* sp. *Biotechnol Lett* 36:341–347
- Herai S, Hashimoto Y, Higashibata H, Maseda H, Ikeda H, Omura S, Kobayashi M (2004) Hyper-inducible expression system for *Streptomyces*. *Proc Natl Acad Sci* 101:14031–14035
- Kaul P, Banerjee UC (2008) Predicting enzyme behaviour in nonconventional media: correlating nitrilase function with solvent properties. *J Ind Microbiol Biotechnol* 35:713–720
- Kinfe HH, Chhiba V, Frederick J, Bode ML, Mathiba K, Steenkamp PA, Brady D (2009) Enantioselective hydrolysis of β -hydroxy nitriles using the whole cell biocatalyst *Rhodococcus rhodochrous* ATCC BAA-870. *J Mol Catal B Enzym* 59:231–236
- Kobayashi M, Nagasawa T, Yamada H (1989) Nitrilase of *Rhodococcus rhodochrous* J1. *Eur J Biochem* 182:349–356
- Kobayashi M, Goda M, Shimizu S (1998) The catalytic mechanism of amidase also involves nitrile hydrolysis. *FEBS Lett* 439:325–328
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680–685
- Layh N, Parratt J, Willetts A (1998) Characterization and partial purification of an enantioselective arylacetone nitrilase from *Pseudomonas fluorescens* DSM 7155. *J Mol Catal B Enzym* 5:476–474
- Leonova TE, Astaurova OB, Ryabchenko LE, Yanenko AS (2000) Nitrile hydratase of *Rhodococcus*: optimization of synthesis in cells and industrial applications for acrylamide production. *Appl Biochem Biotechnol* 88:231–241
- Lévi-Schil S, Soubrier F, Crutz-Le Coq A-M, Faucher D, Crouzet J, Pétre D (1995) Aliphatic nitrilase from a soil-isolated *Comomonas testosteroni* sp.: gene cloning and overexpression, purification and primary structure. *Gene* 161:15–20
- Martínková L, Uhnáková B, Pátek M, Nešvera J, Křen V (2009) Biodegradation potential of the genus *Rhodococcus*. *Environ Int* 35:162–177
- Martínková L, Rucká L, Nešvera J, Pátek M (2017) Recent advances and challenges in the heterologous production of microbial nitrilases for biocatalytic applications. *World J Microbiol Biotechnol* 33:8
- Mateo C, Chmura A, Rustler S, van Rantwijk F, Stolz A, Sheldon RA (2006) Synthesis of enantiomerically pure (S)-mandelic acid using an oxynitrilase-nitrilase bienzymatic cascade: a nitrilase surprisingly shows nitrile hydratase activity. *Tetrahedron Asymmetry* 17:320–323
- Miller JM, Conn EE (1980) Metabolism of hydrogen cyanide by higher plants. *Physiol Plant* 65:1199–1202
- Nagasawa T, Ryuno K, Yamada H (1986) Nitrile hydratase of *Brevibacterium* R-312 purification and characterisation. *Biochem Biophys Res Commun* 139:1304–1312
- Nagasawa T, Nakamura T, Yamada H (1990) ϵ -Caprolactam, a new powerful inducer for the formation of *Rhodococcus rhodochrous* J1 nitrilase. *Arch Microbiol* 155:13–17
- Nagasawa T, Takeuchi K, Yamada H (1991) Characterization of a new cobalt-containing nitrile hydratase purified from urea-induced cells of *Rhodococcus rhodochrous* J1. *Eur J Biochem* 196:581–589
- Nagasawa T, Weiser M, Nakamura T, Iwahara H, Yoshida T, Gekko K (2000) Nitrilase or *Rhodococcus rhodochrous* J1. Conversion into the active form by subunit association. *Eur J Biochem* 267:138–144
- Nageshwar YVD, Sheelu G, Shambu RR, Muluka H, Mehdi N, Malik MS, Kamal A (2011) Optimization of nitrilase production from *Alcaligenes faecalis* MTCC 10757 (ICT-A3): effect of inducers on substrate specificity. *Bioprocess Biosyst Eng* 34:515–523
- Pertsovich SI, Guranda DT, Podchernyaev DA, Yanenko AS, Svedas VK (2005) Aliphatic amidase from *Rhodococcus rhodochrous* M8 is related to the nitrilase/cyanide hydratase family. *Biochem Mosc* 70:1280–1287
- Petříčková A, Veselá AB, Kaplan O, Kubáč D, Uhnáková B, Malandra A, Felsberg J, Rinágelová A, Weyrauch P, Křen V, Bezouška K, Martínková L (2012) Purification and characterization of heterologously expressed nitrilases from filamentous fungi. *Appl Microbiol Biotechnol* 93:1553–1561
- Prasad S, Misra A, Jangir VP, Awasthi A, Raj J, Bhalla TC (2007) A propionitrile-induced nitrilase of *Rhodococcus* sp. NDB 1165 and its application in nicotinic acid synthesis. *World J Microbiol Biotechnol* 23:345–353
- Prasad S, Raj J, Bhalla TC (2009) Purification of a hyperactive nitrile hydratase from resting cells of *Rhodococcus rhodochrous* PA-34. *Indian J Microbiol* 49:237–242
- Precigou S, Wieser M, Pommarens P, Goulas P, Duran R (2004) *Rhodococcus pyridinovorans* MW3, a bacterium producing a nitrile hydratase. *Biotechnol Lett* 26:1379–1384
- Rapheeha OK, Roux-van der Merwe MP, Badenhorst J, Chhiba V, Bode ML, Mathiba K, Brady D (2017) Hydrolysis of nitriles by soil bacteria: variation with soil origin. *J Appl Microbiol* 122:686–697
- Robinson WG, Hook RH (1964) Ricinine nitrilase: I. Reaction product and substrate specificity. *J Biol Chem* 239:4257–4262
- Sharma NN, Sharma M, Bhalla TC (2011) An improved nitrilase-mediated bioprocess for synthesis of nicotinic acid from 3-cyanopyridine with hyperinduced *Nocardia globerula* NHB-2. *J Ind Microbiol Biotechnol* 38:1235–1243
- Stalker DM, Maljy LB, McBride KE (1988) Properties of a nitrilase specific for the herbicide bromoxynil and corresponding nucleotide sequence analysis of the *bnx* gene. *J Biol Chem* 263:6310–6314
- Stevenson DE, Feng R, Dumas F, Groleau D, Mihoc A, Storer AC (1992) Mechanistic and structural studies on *Rhodococcus* ATCC 39484 nitrilase. *Biotechnol Appl Biochem* 15:283–302
- Tani Y, Kurihara M, Nishise H (1989) Characterization of nitrile hydratase and amidase, which are responsible for the conversion of dinitriles to mononitriles, from *Corynebacterium* sp. *Agric Biol Chem* 53:3151–3158
- Tauber MM, Cavaco-Paulo A, Robra K, Gubitz GM (2000) Nitrile hydratase and amidase from *Rhodococcus rhodochrous* hydrolyze

- acrylic fibers and granular polyacrylonitriles. *Appl Environ Microbiol* 66:1634–1638
- Thalenfeld B, Grossowicz N (1976) Regulatory properties of an inducible aliphatic amidase in a thermophilic *Bacillus*. *J Gen Microbiol* 94: 131–141
- Thuku RN, Brady D, Benedik MJ, Sewell BT (2009) Microbial nitrilases: versatile, spiral forming enzymes. *J Appl Microbiol* 106:703–727
- Vaudel M, Burkhardt JM, Zahedi RP, Oveland E, Berven FS, Sickmann A, Martens L, Barsnes H (2015) PeptideShaker enables reanalysis of MS-derived proteomics data sets. *Nat Biotechnol* 33:22–24
- Vejvoda V, Kubác D, Davidová A, Kaplan O, Šulc M, Šveda O, Chaloupková R, Martinkova L (2010) Purification and characterization of nitrilase from *Fusarium solani* IMI96840. *Process Biochem* 45:1115–1120
- Yamada H, Kobayashi M (1996) Nitrile hydratase and its application to industrial production of acrylamide. *Biosci Biotechnol Biochem* 60: 1391–1400
- Yusuf F, Chaubey A, Jamwal U, Parshad R (2013) A new isolate from *Fusarium proliferatum* (AUF-2) for efficient nitrilase production. *Appl Biochem Biotechnol* 171:1022–1031
- Zangi R, Zhou R, Berne BJ (2009) Urea's action on hydrophobic interactions. *J Am Chem Soc* 131:1535–1541
- Zhang ZJ, Xu JH, He YC, Ouyang LM, Liu YY (2011) Cloning and biochemical properties of a highly thermostable and enantioselective nitrilase from *Alcaligenes* sp. ECU0401 and its potential for (R)-(-)-mandelic acid production. *Bioprocess Biosyst Eng* 34:315–322
- Zheng JG, Chen J, Liu ZQ, Wu MH, Xing LY, Shen YC (2008) Isolation, identification and characterization of *Bacillus subtilis* ZJB-063, a versatile nitrile-converting bacterium. *Appl Microbiol Biotechnol* 77:985–993

Websites

Merops-the peptidase database 2017, accessed 20th March 2017, <<https://www.ebi.ac.uk/merops/cgi-bin/speccards?sp=sp034913;type=peptidase>>

Reproduced with permission of copyright owner. Further reproduction prohibited without permission.