

**PREVALENCE, CAUSATIVE FACTORS AND MANAGEMENT OF NOCTURNAL
ENURESIS IN SOUTH AFRICAN CHILDREN**

Margaret Willemina Fockema

**Dissertation submitted to the Faculty of Health Sciences, University of the
Witwatersrand, Johannesburg in fulfillment of the requirements for the degree**

of

MSc Med in Urology

Johannesburg, 2008

DECLARATION

I, Margaret Willemina Fockema declare that this dissertation is my own work. It is being submitted for the degree of Master of Medicine in the branch of Urology in the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at this or any other University.

..... (Signature of Candidate)

.....day of.....2008

DEDICATION

I dedicate this work to my daughters Jeanette and Elisca

Publications and presentations

Fockema MW. (2006) Enuresis in South Africa. All for Kids Congress 2006

Abstract

Prevalence, causative factors and management of nocturnal enuresis in South African children

Objective

A cross sectional study using a self-administered questionnaire to establish the prevalence and causative factors of nocturnal enuresis in 5-10 year old South African children.

Subjects and methods

A questionnaire was distributed to the parents of 4700 school going children between the ages of 5-10 years in selected areas of Gauteng, Kwa-Zulu Natal and the Western Cape in South Africa. Where permission was granted, schools listed in these areas, and schools from different socio-economic areas (suburbs differing in affluence), were contacted telephonically. Those willing to participate were included in the study. Parents anonymously filled out the questionnaire and returned it to school. Data were reported as frequency and percentages in tables and graphs. The Chi-square test was used to compare proportions between groups with Fisher's Exact test used to correct for small numbers of observations ($n \leq 5$). Age differences were determined using Student's t-test. A p value of less than 0.05 was considered to be statistically significant.

Results

The questionnaire response rate was 72.1%. The overall prevalence of nocturnal enuresis was 16.0%, 14.4% for mono-symptomatic nocturnal enuresis and 1.6% for daytime incontinence. A positive family history was found in 50.5% children and constipation was a problem in 16.0% of children with enuresis. Various stress factors were linked to enuresis according to

the parents' perceptions. Only 28.3% had received a form of treatment and 13.5% had been treated by a doctor. A higher prevalence rate of enuresis could not be linked to a lower socio-economic grouping. Parents' awareness of treatment modalities available to their children is outdated and most of the management of the problem was done by the parents themselves.

Conclusion

These results suggest that the prevalence of nocturnal enuresis and associated causative factors in South Africa is similar to the prevalence in other countries, although it is difficult to compare these prevalence rates as studies use different inclusion criteria.

Acknowledgements

- 1. My supervisor: Professor M Haffejee for his help and encouragement.**
- 2. Dr G. Candy for all his advice, assistance and encouragement.**
- 3. My parents for their help and support.**
- 4. My children for their help and support and patience.**
- 5. Ferring Pharmaceuticals for their financial support and support in distributing the questionnaires.**

TABLE OF CONTENTS

	Page
DECLARATION	ii
DEDICATION	iii
PUBLICATIONS AND PRESENTATIONS	iv
ABSTRACT	v
ACKNOWLEDGEMENTS	vii
TABLE OF CONTENTS	viii
LIST OF FIGURES	xii
LIST OF TABLES	xiii
NOMENCLATURE	xv
1.0 INTRODUCTION	1
1.1 Literature review	1
1.1.1 Definition	1
1.1.2 Prevalence	4
1.1.3 Severity	6
1.1.4 Gender differences	6
1.1.5 Heredity	7
1.1.6 Low self esteem, psychological problems and poor school performance	7
1.1.7 Socio-economic factors	8
1.1.8 Stressful events	10

1.1.9 Other associated factors	10
1.1.10 Perceptions of parents	10
1.2 Summary	12
2.0 METHODS	13
2.1 Study questionnaire	13
2.1.1 Sample size calculation	14
2.1.2 Selection of study sites	15
2.2 Procedure	15
2.2 .1 Data recording and statistical analysis	16
2.2.2 Socio-economic status	17
2.2.3 Statistical analysis	17
2.2.4 Study limitations	17
3.0 RESULTS	19
3.1 Prevalence	19
3.1.1 Overall prevalence of MNE and PNE in different age groups	19
3.1.2 Overall prevalence of MNE according to province	21
3.1.2.1 Prevalence of children with MNE in the Western Cape	23
3.1.2.2 Prevalence of children with MNE in Kwa-Zulu Natal	23
3.1.2.3 Prevalence of children with MNE in Gauteng	24

3.1.3 Prevalence of daytime urinary incontinence	25
3.1.4 Prevalence according to gender	26
3.2 Frequency of NE	27
3.3 Potential risk factors for NE	29
3.3.1 Heredity	29
3.3.2 Constipation	30
3.3.3 Possible link to stress	31
3.3.4 Performance at school	33
3.4 Awareness of treatment	34
3.4.1 Number and percentage of children that parents would take for treatment	35
3.4.2 Number and percentage of children treated for NE	36
3.4.3 Treatment modalities	38
3.4.4 Medicine given	39
3.5 Socio-economic status	40
4.0 DISCUSSION	41
4.1 Study outcome	41
4.2 Prevalence of NE	42
4.2.1 Prevalence of daytime urinary incontinence	44
4.2.2 Prevalence according to gender	44
4.3 Frequency of NE	44
4.4 Potential risk factors for NE	46

4.4.1 Heredity	46
4.4.2 Constipation	47
4.4.3 Possible link to stress	48
4.4.3.1 Possible stressful events	49
4.4.4 Performance at school	49
4.4.5 Socio-economic status	50
4.4.6 Ethnicity	50
4.5 Awareness of treatment	51
4.5.1 Percentage of children that parents would take for treatment	51
4.5.2 Percentage of children treated for NE	51
4.5.3 Treatment modalities	52
4.5.4 Medicine given	54
5.0 CONCLUSIONS	55
6.0 RECOMMENDATIONS	57
6.1 Awareness about NE	57
6.2 Future studies about NE	58
ANNEXURE 1	59
ANNEXURE 2	63
REFERENCES	64

LIST OF FIGURES

Pages

Figure

1.1 Percentage of study subjects with NE who visited a health care worker	11
3.1 Overall prevalence of MNE in South Africa	19
3.2 Overall prevalence rates in different provinces	22
3.3 Percentage of enuretic children with stress factors	32
3.4 Parent's awareness of available treatment as a percentage	34
3.5 Number of children receiving treatment (from Table 3.15)	37
4.1 Comparison of frequency of NE between ALSPAC study and the present study	45
4.2 Prevalence as a percentage of familial NE in Bangkok	47

LIST OF TABLES

Tables	Page
1.1 Definitions of NE	2
1.2 The prevalence of NE at 7 years old of age (adapted from reference 3)	3
1.3 Prevalence of NE according to age (adapted from reference 5)	5
1.4 Prevalence rates from various regions and socio-economic groups	9
3.1 Overall prevalence of NE according to age	20
3.2 Prevalence rates of MNE in different provinces	21
3.3 Prevalence of MNE in the Western Cape according to age	23
3.4 Prevalence of MNE in Kwa-Zulu Natal according to age	23
3.5 Prevalence of MNE in Gauteng according to age	24
3.6 Prevalence of daytime urinary incontinence	25
3.7 Prevalence according to gender	26
3.8 Frequency of NE episodes according to age	27
3.9 Frequency of NE episodes according to various definitions	28
3.10 Number and percentage of parents and siblings of study subjects with MNE	29
3.11 Children with MNE suffering from constipation	30
3.12 Possible links to stress	31
3.13 Performance at school	33

Tables cont	Page
3.14 Percentage of parents who would take children for treatment	35
3.15 Children with NE receiving treatment	36
3.16 Treatment modalities chosen by parents of children with NE	38
3.17 Types of medicine given to treat NE	39
3.18 Socio-economic status* in subjects with NE	40
4.1 Prevalence of NE among 7 year olds where ICD 10 definition was used	43
4.2 Parental strategies for managing NE in different countries	53

Nomenclature

Abbreviations (Acronyms)

NE	Nocturnal enuresis
PNE	Poly-symptomatic nocturnal enuresis
MNE	Mono-symptomatic nocturnal enuresis
CNS	Central Nervous system
DSM III	Diagnostic and Statistical Manual of Mental disorders, Third Edition
DSM IV	Diagnostic and Statistical Manual of Mental disorders, Fourth Edition
ICD-10	International Classification of Diseases and Related Health Problems 10 th revision version for 2007
ICCS	International Children's Continence Society
ICS	International Continence Society