

Hearing Function in Children with Chronic Renal Dysfunction

Student: Jennifer Lau

Submitted to the

Discipline of Speech Pathology and Audiology

School of Human and Community Development

Faculty of Humanities

University of the Witwatersrand

in fulfilment of the requirements for the

Degree of Master of Audiology

July 2012

Declaration

I, Jennifer Lau, hereby declare that this submission is my own original work and that the assistance which I have received is detailed in the Acknowledgements of this report. To the best of my knowledge and belief, it contains no material which has been accepted for the award of any other degree or diploma at any other university or other institute of higher learning, except where due acknowledgement has been made in the text. I am responsible for the study and conclusions reached.

JENNIFER LAU

Date

Acknowledgements

I would like to express my sincerest gratitude to the following people:

- ❖ Professor Katijah Khoza-Shangase, my supervisor, for her encouragement, advice and assistance throughout the supervision of my research.
- ❖ The participants with chronic renal disorder and their caregivers, who so willingly gave of their time to expand my knowledge.
- ❖ The Chief Executive Officer, Medical Superintendent and Heads of Department for both the Renal Department and Speech and Hearing Department at Charlotte Maxeke Johannesburg Academic Hospital, for granting me the opportunity to conduct this research.
- ❖ Dr. Levy, for his time and knowledge that he shared with me regarding renal disorders as well as the advice that he gave me regarding the research. I would also like to thank all the other doctors, nurses and administrative staff from the Renal Department, who so willingly helped me during the data collection phase of the study.
- ❖ The Head of Department as well as my fellow friends and colleagues from the Speech and Hearing Department, for their continual support, friendship, encouragement and assistance during my academic endeavours.
- ❖ The Ear, Nose and Throat Specialists at Charlotte Maxeke Johannesburg Academic Hospital, for their support and encouragement.
- ❖ Professor Peter Fridjhon, Dr. Michael Greyling, and Dr. Charles Chimedza, statisticians from the University of the Witwatersrand, for all their valuable time, knowledge and guidance regarding data analysis of the study. Their input and guidance is much appreciated.
- ❖ Mr. Herman Tesner, for his most timeous input, guidance and support with regards to the final check and editing of the research report.
- ❖ All hospital speech and hearing therapists and university staff who provided audiological input and emotional encouragement during this study.
- ❖ My family, for their unconditional love and support in the realization of my dreams, and standing by me through thick and thin.
- ❖ My friends, for their encouragement, interest and patience in the carrying out of my research study.

May God Bless You Always!

Abstract

The primary aim of the research was to describe hearing function in a group of children with chronic renal dysfunction receiving treatment in an academic hospital in Johannesburg, South Africa. Specific objectives in the study were to determine the prevalence of hearing loss in paediatric patients with chronic renal dysfunction; to describe the type, degree and configuration of the hearing loss; and to establish if there was a relationship between the presenting hearing loss and the severity of renal dysfunction, the different treatment regimens, duration of renal dysfunction, and the duration of treatment.

One hundred children between the ages five -18 years participated in the study and comprised 65 males and 35 females. The mean age of the participants was 11.68 years.

A cross-sectional, descriptive, quantitative research design was employed. All participants underwent a case history interview and a full audiological examination which included an otoscopic examination, immittance testing (tympanometry and ipsilateral acoustic reflex testing), pure tone audiometry including extended high frequency testing up to 16 kilohertz as well as diagnostic distortion product otoacoustic emission testing. A record review was also done.

Both descriptive and inferential statistics were used to analyse the collected data. Inferential statistics included parametric measures using multiple regression measures as well as non parametric measures using the Kruskal-Wallis statistical analysis.

Results revealed that there was a high prevalence of hearing loss in children with chronic renal dysfunction. Results from the extended high frequency pure tone testing as well as the diagnostic distortion product testing revealed that the most common hearing loss was a low and high to ultrahigh frequency mild sensorineural hearing loss. The study showed that there was no relationship between the severity of hearing loss and the severity of renal dysfunction, or the duration of renal dysfunction and the duration of treatment. However, the study showed that there was a relationship between the severity of hearing loss and certain treatments, that is,

haemodialysis and the use of ototoxic medication such as loop diuretics, tuberculosis medication, and antimalarial medication.

As the potential to miss hearing loss in this population is high, the research highlighted the importance of extended high frequency audiometry as well as diagnostic distortion product otoacoustic emission testing for the use of ototoxic monitoring in patients with chronic renal dysfunction. The research also highlighted the need for further research in this area as well as the need for educating medical personnel and caregivers working with children with chronic renal disease.

Key words: chronic renal dysfunction; hearing loss; distortion product; otoacoustic emissions; extended high frequency audiometry.

Table of Contents

	Pages
Acknowledgements	iii
Abstract	iv-v
List of Tables	viii-ix
List of Figures	x
List of Appendices	xi
Introduction	1-22
Methodology	23-40
Aims	23
Research Design	24
Participants	24-25
Research instrumentation	26-27
Ethical considerations	27-28
Research procedure	28-30
Audiological testing	30-37
Validity and reliability	37
Analysis of data	37-40
Results	41-74
Description of participants	41-42
Data analysis in terms of case history	43-46
Description of hearing function	47-56
Prevalence of hearing loss	56-57
Type, degree and configuration of hearing loss	57-61
Relationship between hearing loss and severity of chronic renal disease	61-63
Relationship between hearing loss and the different treatment regimens	64-66
Relationship between hearing loss and the duration of renal dysfunction	66-67
Relationship between hearing loss and the duration of treatment for renal dysfunction	68-70
Additional results obtained of the study	70-74
Discussion of results	75-82
Conclusion, limitations and recommendations	83-88

References
Appendices

89-97
98-113

List of tables

	Pages
Table 1: Stages of Chronic Kidney Disease	2
Table 2: Prevalence Figures of Hearing Loss for Sub-Saharan Africa	12
Table 3: Published Studies on the Effect of Haemodialysis on Hearing Function	16-17
Table 4: Results of Loop Diuretic Effects on Hearing Function in Animal Studies	20-21
Table 5: Interpretation of Tympanometry Results	32
Table 6: Classification of Hearing Loss	34-35
Table 7: Mean Pure Tone Audiometry Results, Standard Deviations and Median Results for the Right and Left Ear (in dB HL)	51
Table 8: Pure Tone Audiometry Results for the Right and Left Ear	52-53
Table 9: Mean DPOAE Results and Standard Deviations for the Right and Left Ear	54-55
Table 10: Type and degree of hearing loss between ears	60-70
Table 11: Parametric analysis showing the relationship between severity of hearing loss and stage of chronic renal dysfunction	62
Table 12: Non-parametric statistical analysis showing the relationship between degree of hearing loss and severity of chronic renal dysfunction	63
Table 13: Parametric Statistical Analysis of the independency of variable severity of hearing loss and treatment methods	64
Table 14: Non-Parametric Statistical Analysis of the relationship between degree of hearing loss and different ototoxic medications	65
Table 15: Parametric logistic regression statistical analysis of the relationship between degree of hearing loss and the duration of chronic renal dysfunction	66
Table 16: Non-parametric Kruskal-Wallis statistical analysis of the relationship between degree of hearing loss and the duration of chronic renal dysfunction	67

Table 17:	Parametric logistic regression statistical analysis of the relationship between degree of hearing loss and the duration of treatment for chronic renal dysfunction	68
Table 18:	Non-parametric Kruskal-Wallis statistical analysis of the relationship between degree of hearing loss and duration of treatment for chronic renal dysfunction	69
Table 19:	Parametric multiple regression statistical analysis of the relationship between degree of hearing loss, duration of renal dysfunction and duration of treatment	70

List of figures

	Page
Figure 1: Gender distribution of participants with chronic renal dysfunction	41
Figure 1: Distribution of stages of chronic kidney disease	42
Figure 2: Number of participants with a family history of kidney disease	43
Figure 3: Treatment regimens for participants with chronic renal disorder	44
Figure 4: Medications used by the participants	45
Figure 5: Otoloscopic examination for each participant	47
Figure 6: Tympanometry results for each participant	48
Figure 7: Ipsilateral acoustic reflexes for the right and left ear	49
Figure 8: Mean pure tone audiometry results for the right and left ear for each frequency (in dBHL)	50
Figure 9: Mean DPOAE results for the right and left ear for each frequency	54
Figure 10: DPOAE responses for the right and left ear	56
Figure 11: Prevalence of hearing loss in a sample of paediatric participants with chronic kidney disease	57
Figure 12: Types of hearing loss	58
Figure 13: Degree of hearing loss	59
Figure 14: Perceived hearing loss by the participants and their caregivers	71
Figure 15: Perceived onset of hearing loss	72
Figure 16: Perceived symmetry of hearing loss	73
Figure 17: Other related ear symptoms	74

List of appendices

	Page
Appendix A: Participant information sheet for parents/legal guardians	98-99
Appendix B: Participant information sheet for paediatric patients	100-101
Appendix C: Consent form for parent/legal guardian of the participants in the study	102
Appendix D: Assent form for the participants in the study	103
Appendix E: Hospital information sheet	104-105
Appendix F: Case history questionnaire for parents/legal guardians of paediatric patients with renal dysfunction (Biographical Information)	106
Appendix G: Case history questionnaire for parents/legal guardians of paediatric patients with renal dysfunction (Renal and Hearing Information)	107
Appendix H: Letter of permission for the study from the chief executive officer of Charlotte Maxeke Johannesburg Academic Hospital	108
Appendix I: Letter of permission for the study from the head of the paediatric renal department of Charlotte Maxeke Johannesburg Academic Hospital	109
Appendix J: Letter of permission for the study from the head of the speech and hearing department of Charlotte Maxeke Johannesburg Academic Hospital	110
Appendix K: Letter of permission for the study from Human Research Ethical Committee	111
Appendix L: Certificate of calibration for the GSI Tympanometer	112
Appendix M: Certificate of calibration for the Audera distortion product otoacoustic emission meter	113