



Fejér–Riesz factorization in the QRC-subalgebra and circularity of the quaternionic numerical range

Alma van der Merwe¹ · Madelein van Straaten² · Hugo J. Woerdeman³ 

Received: 23 November 2023 / Accepted: 21 February 2024
© The Author(s) 2024

Abstract

We provide a characterization when the quaternionic numerical range of a matrix is a closed ball with center 0. The proof makes use of Fejér–Riesz factorization of matrix-valued trigonometric polynomials within the algebra of complex matrices associated with quaternion matrices.

Keywords Fejér–Riesz factorization · Quaternionic numerical range · Circularity · QRC-subalgebra

Mathematics Subject Classification 15A60 · 15A63 · 15B33 · 47A12

1 Introduction

In this paper, we are interested in studying when the quaternionic numerical range is the closed unit ball in the algebra of quaternions. The quaternionic numerical range was introduced in [14] and several basic properties were derived in [2, 3, 14, 19]. More recent papers include [1, 6, 17]. For the case of nilpotent quaternionic matrices,

Dedicated to Chi-Kwong Li, a wonderful colleague and a great friend.

Communicated by Douglas Farenick.

✉ Hugo J. Woerdeman
hugo@math.drexel.edu

Alma van der Merwe
alma.vandermerwe@wits.ac.za

Madelein van Straaten
Madelein.VanStraaten@nwu.ac.za

- ¹ Department of Mathematics, University of the Witwatersrand, Johannesburg, South Africa
- ² Department of Mathematics and Applied Mathematics, Research Focus: Pure and Applied Analytics, North-West University, Potchefstroom, South Africa
- ³ Department of Mathematics, Drexel University, 3141, Chestnut Street, Philadelphia, PA 19104, USA

some circularity results appear in [5]. In the case of complex matrices, it was derived (beyond the nilpotent case) in [16, Theorem 4.5] that for a complex matrix $B \in \mathbb{C}^{n \times n}$, its numerical range is the closed unit disk if and only if there exist $P_0, P_1 \in \mathbb{C}^{(n-1) \times n}$, so that $B = 2P_0^*P_1$ and $P_0^*P_0 + P_1^*P_1 = I_n$. We will be deriving a quaternionic version of this result. To do this, we need to study Fejér–Riesz factorization in the algebra of complex matrices associated with quaternion matrices. This algebra, which we will call the *QRC-subalgebra*, consists of matrices of the form

$$\begin{bmatrix} A & \overline{B} \\ -B & A \end{bmatrix} \in \mathbb{C}^{2n \times 2n},$$

where $\overline{C} = (\overline{c_{ij}})$ denotes the matrix obtained from $C = (c_{ij})$ by taking entrywise complex conjugates. Here, ‘QRC’ stands for ‘Quaternions Represented as Complexes’. It should be noted that in [9], for instance, these matrices are referred to as being ‘symplectic’. However, as the term ‘symplectic’ is also used for other types of matrices, we will use the QRC-subalgebra terminology. The classical Fejér–Riesz factorization result has its application in filter design (see, e.g., [11]), wavelet design (see [7]), and H_∞ -control (see, e.g., [10]). Factorizations in this context are numerically found via solving Riccati equations (see [15]) or semidefinite programming (see, e.g., [12]).

The paper is organized as follows. In Sect. 2, we will derive the Fejér–Riesz factorization in the QRC-subalgebra. In Sect. 3, we will provide a characterization when the quaternionic numerical range equals the closed unit ball in the quaternions. Before we start, we will introduce our notational conventions.

1.1 Notation

For notation regarding basic quaternion linear algebra, we refer to the monograph [18]. Every quaternion in \mathbb{H} is of the form: quaternion number

$$x = x_0 + x_1i + x_2j + x_3k,$$

where $x_0, x_1, x_2, x_3 \in \mathbb{R}$ and the elements i, j, k satisfy the following formulas:

$$i^2 = j^2 = k^2 = -1, \quad ij = -ji = k, \quad jk = -kj = i, \quad ki = -ik = j.$$

Note that multiplication in \mathbb{H} is not commutative. Denote the *conjugate quaternion* conjugate quaternion of x by $\overline{x} = x_0 - x_1i - x_2j - x_3k$ and the *norm* of $x \in \mathbb{H}$ by $\|x\| = \sqrt{x^*x} = \sqrt{x_0^2 + x_1^2 + x_2^2 + x_3^2} \in \mathbb{R}$. For a quaternion matrix A , let \overline{A} denote the matrix obtained from where each entry is the conjugate of the corresponding entry in A .

For $A \in \mathbb{H}^{n \times n}$, where

$$A = A_0 + iA_1 + jA_2 + kA_3, \quad \text{with } A_i \in \mathbb{R}^{n \times n}, i = 0, 1, 2, 3,$$

we write $A = B + jC$ where $B := A_0 + iA_1 \in \mathbb{C}^{n \times n}$ and $C := A_2 - iA_3 \in \mathbb{C}^{n \times n}$. There exists a complex matrix $\omega_n(A)$ associated with A , defined by

$$\omega_n(A) = \begin{bmatrix} B & \overline{C} \\ -C & \overline{B} \end{bmatrix} \in \mathbb{C}^{2n \times 2n}.$$

Then, ω_n is an isomorphism of the real algebra $\mathbb{H}^{n \times n}$ onto the real unital subalgebra Ω_{2n} (subalgebra of $\mathbb{C}^{2n \times 2n}$)

$$\Omega_{2n} := \left\{ \begin{bmatrix} B & \overline{C} \\ -C & \overline{B} \end{bmatrix} : B, C \in \mathbb{C}^{n \times n} \right\}$$

of $\mathbb{C}^{2n \times 2n}$. As mentioned before, we will call this subalgebra the *QRC-subalgebra* where ‘QRC’ stands for ‘Quaternions Represented as Complexes’. For a nonsquare matrix $A \in \mathbb{H}^{m \times n}$ written as $A = B + jC$, $B, C \in \mathbb{C}^{m \times n}$, we use the notation $\omega_{m,n}(A)$ and $\Omega_{2m,2n}$

$$\Omega_{2m,2n} := \left\{ \begin{bmatrix} B & \overline{C} \\ -C & \overline{B} \end{bmatrix} : B, C \in \mathbb{C}^{m \times n} \right\}.$$

The map $\omega_{m,n} : \mathbb{H}^{m \times n} \rightarrow \Omega_{2m,2n}$ is invertible, and the inverse of $\omega_{m,n}$ is denoted by $\omega_{m,n}^{-1}$. When the sizes are clear from the context, we may drop the subscripts and simply write $\omega(A)$.

The following properties can be found in [18, Section 3.4] and [21]. For $X, Y \in \mathbb{H}^{n \times n}$ and $s, t \in \mathbb{R}$ be arbitrary, we have

- (i) $\omega_n(I_n) = I_{2n}$;
- (ii) $\omega_n(XY) = \omega_n(X)\omega_n(Y)$;
- (iii) $\omega_n(sX + tY) = s\omega_n(X) + t\omega_n(Y)$;
- (iv) $\omega_n(X^*) = (\omega_n(X))^*$;
- (v) X is invertible if and only if $\omega_n(X)$ is invertible; if so, then $\omega_n(X^{-1}) = (\omega_n(X))^{-1}$.

Furthermore, define the matrix

$$J_n := \omega(j) \otimes I_n = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix},$$

where \otimes is the Kronecker product.

We denote the numerical range of a quaternion matrix $A \in \mathbb{H}^{n \times n}$ by

$$W_{\mathbb{H}}(A) = \{x^*Ax : \|x\| = 1, x \in \mathbb{H}^n\} \subset \mathbb{H}.$$

Note that if $q \in W_{\mathbb{H}}(A)$, and $u \in \mathbb{H}$ where $\|u\| = 1$, then $u^*qu \in W_{\mathbb{H}}(A)$. In general, $u^*qu \neq q$. For basic information on the quaternionic numerical range, see for instance

[18, Section 3.3]. The subset of complex elements in $W_{\mathbb{H}}(A)$ is known as the *bild* of A , denoted

$$B(A) := W_{\mathbb{H}}(A) \cap \mathbb{C}.$$

Although the bild may not be convex, the upper bild $B^+(A) = W_{\mathbb{H}}(A) \cap \mathbb{C}^+$ is always convex; see [19]. Here, $\mathbb{C}^+ = \{z \in \mathbb{C} : \text{Im}z \geq 0\}$ is the closed upper half plane. Note that $B(A)$ is symmetric with respect to the real axis. Indeed, if $j^* \alpha j = \bar{\alpha} \in B(A)$. Thus, knowing just $B^+(A)$ provides full information on $B(A)$, since $B(A) = B^+(A) \cup \overline{B^+(A)}$ (where the bar indicates taking complex conjugates).

Let $\mathbb{D} = \{z \in \mathbb{C} : \|z\| < 1\}$ and $\overline{\mathbb{D}} = \{z \in \mathbb{C} : \|z\| \leq 1\}$ denote the open and closed unit disks in \mathbb{C} and $\mathbb{T} = \{z \in \mathbb{C} : \|z\| = 1\}$ the unit circle. Finally, let $\mathbb{B} = \{q \in \mathbb{H} : \|q\| < 1\}$ and $\overline{\mathbb{B}} = \{q \in \mathbb{H} : \|q\| \leq 1\}$ denote the open and closed unit balls in \mathbb{H} .

2 Fejér–Riesz factorization in the QRC-subalgebra

Lemma 2.1 *Let $C \in \mathbb{C}^{2n \times 2n}$. Then $C \in \Omega_{2n}$ if and only if $J_n \overline{C} J_n^* = C$.*

Proof Take $C \in \mathbb{C}^{2n \times 2n}$, where

$$C = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}.$$

Then

$$J_n \overline{C} J_n^* = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} \begin{bmatrix} \overline{C}_{11} & \overline{C}_{12} \\ \overline{C}_{21} & \overline{C}_{22} \end{bmatrix} \begin{bmatrix} 0 & -I_n \\ I_n & 0 \end{bmatrix} = \begin{bmatrix} \overline{C}_{22} & -\overline{C}_{21} \\ -\overline{C}_{12} & \overline{C}_{11} \end{bmatrix},$$

which is equal to C if and only if $C_{11} = \overline{C}_{22}$ and $C_{12} = -\overline{C}_{21}$. Hence, $C \in \Omega_{2n}$ if and only if $J_n \overline{C} J_n^* = C$. □

Lemma 2.2 *Let $C \in \mathbb{C}^{2n \times 2n}$. Then, $\frac{1}{2} (J_n \overline{C} J_n^* + C) \in \Omega_{2n}$.*

Proof Take $C \in \mathbb{C}^{2n \times 2n}$, where

$$C = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}.$$

Then

$$\begin{aligned} \frac{1}{2} (J_n \overline{C} J_n^* + C) &= \frac{1}{2} \left(\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} \begin{bmatrix} \overline{C}_{11} & \overline{C}_{12} \\ \overline{C}_{21} & \overline{C}_{22} \end{bmatrix} \begin{bmatrix} 0 & -I_n \\ I_n & 0 \end{bmatrix} + \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} \right) \\ &= \frac{1}{2} \left(\begin{bmatrix} \overline{C}_{22} & -\overline{C}_{21} \\ -\overline{C}_{12} & \overline{C}_{11} \end{bmatrix} + \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} \right) \\ &= \frac{1}{2} \begin{bmatrix} \overline{C}_{22} + C_{11} & -\overline{C}_{21} + C_{12} \\ -\overline{C}_{12} + C_{21} & \overline{C}_{11} + C_{22} \end{bmatrix} \in \Omega_{2n}. \end{aligned}$$
□

Note that if $C \in \Omega_{2n}$, then $J_n \overline{C} J_n^* = C$, and hence

$$\frac{1}{2} (J_n \overline{C} J_n^* + C) = C.$$

The following result provides Fejér–Riesz factorization in the QRC-subalgebra Ω_{2n} . The result is a variation of [12, Proposition 4.6], which was based on [8, Section 3]; see also [4, Theorem 2.4.19].

Theorem 2.3 *Let $Q(z) = Q_{-m} z^{-m} + \dots + Q_m z^m \geq 0, z \in \mathbb{T}$, where $Q_j \in \Omega_{2n}$. Let*

$$Z = \begin{bmatrix} Q_0 & Q_{-1} & \dots & Q_{-m} \\ Q_1 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ Q_m & 0 & \dots & 0 \end{bmatrix}, \text{ and introduce the convex set}$$

$$\mathcal{G} = \left\{ X \in \mathbb{C}^{2nm \times 2nm} \mid A(X) := Z - \begin{bmatrix} X & 0 \\ 0 & 0_{2n} \end{bmatrix} + \begin{bmatrix} 0_{2n} & 0 \\ 0 & X \end{bmatrix} \geq 0 \right\}.$$

Then, \mathcal{G} has elements X_{\max} and X_{\min} that are maximal and minimal with respect to the Loewner ordering, respectively; i.e., $X_{\max}, X_{\min} \in \mathcal{G}$ have the property that $X \in \mathcal{G}$ implies $X_{\min} \leq X \leq X_{\max}$. Writing $X_{\max} = (X_{ij}^{\max})_{i,j=1}^m$ and $X_{\min} = (X_{ij}^{\min})_{i,j=1}^m$, we have that $X_{ij}^{\max}, X_{ij}^{\min} \in \Omega_{2n}$.

Moreover, consider the set

$$\mathcal{A} = \left\{ A = (A_{ij})_{i,j=0}^m \in \mathbb{C}^{2n(m+1) \times 2n(m+1)} \mid A = A(X) \text{ for some } X \in \mathcal{G} \right\}.$$

Then, $A(X_{\max})$ is the unique element in \mathcal{A} , so that A_{mm} is maximal (or equivalently, A_{00} is minimal) in the Loewner order. Also, $A(X_{\min})$ is the unique element in \mathcal{A} so that A_{00} is maximal (or equivalently, A_{mm} is minimal) in the Loewner order. Finally, we may factor $A(X_{\max})$ and $A(X_{\min})$ as

$$A(X_{\max}) = \begin{bmatrix} H_0^* \\ \vdots \\ H_m^* \end{bmatrix} [H_0 \dots H_m], \quad A(X_{\min}) = \begin{bmatrix} K_0^* \\ \vdots \\ K_m^* \end{bmatrix} [K_0 \dots K_m] \quad (2.1)$$

with $H_i, K_i \in \Omega_{2n}, i = 0, \dots, m$, and put $H(z) = \sum_{k=0}^m H_k z^k, K(z) = \sum_{k=0}^m K_k z^k$. Then

$$Q(z) = H(z)^* H(z) = K(z)^* K(z), \quad z \in \mathbb{T}.$$

Proof Without the restriction that $Q_j, H_j, K_j \in \Omega_{2n}$, the statement is covered by [12, Proposition 4.6]. For a block matrix $B = (B_{ij})_{i,j}$, where $B_{ij} \in \mathbb{C}^{2n \times 2n}$, we define

$$\mathcal{J}(B) := \left(\frac{1}{2} (J_n \overline{B_{ij}} J_n^* + B_{ij}) \right)_{i,j}.$$

In other words, we perform the operation $C \mapsto \frac{1}{2} (J_n \bar{C} J_n^* + C)$ to each block entry. It is easy to see that the operation \mathcal{J} preserves positive semidefiniteness. In addition, $\mathcal{J}(Z) = Z$ and $\mathcal{J}(A(X)) = A(\mathcal{J}(X))$. This now yields that when $X \in \mathcal{G}$, then $\mathcal{J}(X) \in \mathcal{G}$. As \mathcal{J} preserves positive semidefiniteness, it also follows that $X \leq Y$ implies $\mathcal{J}(X) \leq \mathcal{J}(Y)$. Thus, $X_{\min} \leq X \leq X_{\max}$ implies $\mathcal{J}(X_{\min}) \leq \mathcal{J}(X) \leq \mathcal{J}(X_{\max})$. As $\mathcal{J}(X_{\min}), \mathcal{J}(X_{\max}) \in \mathcal{G}$, we also have $X_{\min} \leq \mathcal{J}(X_{\min})$ and $\mathcal{J}(X_{\max}) \leq X_{\max}$. Observe that \mathcal{J} does not change the trace of a Hermitian matrix, and thus, $\text{Tr}(X_{\min}) = \text{Tr}(\mathcal{J}(X_{\min}))$ and $\text{Tr}(\mathcal{J}(X_{\max})) = \text{Tr}(X_{\max})$. But then, it follows that $\mathcal{J}(X_{\min}) = X_{\min}$ and $\mathcal{J}(X_{\max}) = X_{\max}$, and thus $X_{ij}^{\max}, X_{ij}^{\min} \in \Omega_{2n}$. Finally, the spectral theorem for quaternionic matrices [9, Theorem 3.3 and Proposition 3.7] implies that the factorizations in (2.1) can be kept within the QRC-subalgebra. \square

In fact, we have that $H(z)$ is co-outer and $K(z)$ is outer (see [12, Section 4] or [4, Section 2.4] for the definitions); when H_m and K_0 are invertible, this means that $\det H(z) \neq 0$ for $z \in \mathbb{C} \setminus \bar{\mathbb{D}}$ and $\det K(z) \neq 0$ for $z \in \mathbb{D}$.

3 Circularity of the quaternionic numerical range

For $A \in \mathbb{H}^{n \times n}$, we consider also the numerical range of the complex matrix $\omega(A)$

$$W(\omega(A)) = \left\{ y^* \omega(A) y : \|y\| = 1, y \in \mathbb{C}^{2n} \right\}.$$

Since $W(\omega(A))$ is always convex by the famous Toeplitz–Hausdorff Theorem [13, 20] and $B(A)$ sometimes not, we know that the two sets do not coincide in general. However, we will see that $B(A)$ is the unit disk if and only if $W(\omega(A))$ is the unit disk.

We start off with some auxiliary results. The following lemma follows from the proof of [14, Theorem 35].

Lemma 3.1 [14] *For $A \in \mathbb{H}^{n \times n}$ follows $B(A) \subseteq W(\omega(A))$.*

Proof Let $q \in B(A)$. Hence, there exists a vector $x \in \mathbb{H}^n$, where $\|x\| = 1$, such that $q = x^* A x \in \mathbb{C}$. Thus, $\omega(q) = \omega(x)^* \omega(A) \omega(x)$, where

$$\omega(q) = \begin{bmatrix} q & 0 \\ 0 & \bar{q} \end{bmatrix} = \begin{bmatrix} x_C^* & -x_H^* \\ \bar{x}_H^* & \bar{x}_C^* \end{bmatrix} \begin{bmatrix} A_C & \bar{A}_H \\ -A_H & \bar{A}_C \end{bmatrix} \begin{bmatrix} x_C & \bar{x}_H \\ -x_H & \bar{x}_C \end{bmatrix},$$

with $A = A_C + jA_H$ and $x = x_C + jx_H$. Using that

$$1 = \|x\|^2 = \|x_C + jx_H\|^2 = \|x_C\|^2 + \|x_H\|^2 = \left\| \begin{bmatrix} x_C \\ -x_H \end{bmatrix} \right\|^2,$$

it follows that

$$q = [x_C^* \ -x_H^*] \omega(A) \begin{bmatrix} x_C \\ -x_H \end{bmatrix} \in W(\omega(A)). \quad \square$$

In general, the other inclusion will not hold. For example, if $A \in \mathbb{H}^{1 \times 1}$ is not real, then $B(A) = \{\alpha, \bar{\alpha}\}$ where α and $\bar{\alpha}$ are the eigenvalues of $\omega(A) \in \mathbb{C}^{2 \times 2}$; and $W(\omega(A))$ will be the line segment between α and $\bar{\alpha}$.

Next, we show that the two sets $B(A)$ and $W(\omega(A))$ do coincide when either one is the closed unit disc, with $B(A) = \overline{\mathbb{D}}$ the stronger condition, since $W(\omega(A))$ is convex.

Lemma 3.2 *For $A \in \mathbb{H}^{n \times n}$, it follows that $B(A) = \overline{\mathbb{D}}$ is equivalent to $W_{\mathbb{H}}(A) = \overline{\mathbb{B}}$.*

Proof Assume $B(A) = \overline{\mathbb{D}}$. Let $q \in \overline{\mathbb{B}}$ where $q = q_0 + iq_1 + jq_2 + kq_3$ with $\|q\|^2 = \sum_{p=0}^3 |q_p|^2 \leq 1$. Then

$$\tilde{q} = q_0 + i \left\| \begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} \right\| \in \overline{\mathbb{D}} = B(A) = W_{\mathbb{H}}(A) \cap \mathbb{C}.$$

Hence, $\tilde{q} \in W_{\mathbb{H}}(A)$ and by [2, Lemma 2] also $q \in W_{\mathbb{H}}(A)$. For the other inclusion, let $q \in W_{\mathbb{H}}(A)$ where $q = q_0 + iq_1 + jq_2 + kq_3$. Then

$$q_0 + i \left\| \begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} \right\| \in B(A) = \overline{\mathbb{D}},$$

from which follows

$$1 \geq |q_0|^2 + \left\| \begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} \right\|^2 = \sum_{p=0}^3 |q_p|^2 = \|q\|^2.$$

Hence, $q \in \overline{\mathbb{B}}$.

For the other direction, assume $W_{\mathbb{H}}(A) = \overline{\mathbb{B}}$. Take $z \in B(A) = W_{\mathbb{H}}(A) \cap \mathbb{C}$. Thus, $\|z\|^2 \leq 1$ which proves $z \in \overline{\mathbb{D}}$. For the other inclusion, take $z \in \overline{\mathbb{D}}$ where $z = x + iy$ and $x, y \in \mathbb{R}$. Thus $\|z\|^2 = x^2 + y^2 \leq 1$. Then, $x + iy \in \mathbb{H}$ and $x + iy \in \overline{\mathbb{B}} = W_{\mathbb{H}}(A)$. Using [2, Lemma 2], it follows that $z \in B(A)$. □

The following lemma follows from [2, Theorem 2], which states that if $W_{\mathbb{H}}(A)$ is convex, then $B(A) = W(\omega(A))$. We will provide a proof as it makes the paper more self-contained.

Lemma 3.3 *For $A \in \mathbb{H}^{n \times n}$, if $B(A) = \overline{\mathbb{D}}$, then $B(A) = W(\omega(A))$.*

Proof If $B(A) = \overline{\mathbb{D}}$, then $\overline{\mathbb{D}} \subseteq W(\omega(A))$ using Lemma 3.1. Hence, it remains to show that $W(\omega(A)) \subseteq \overline{\mathbb{D}}$.

Since $B(A) = \overline{\mathbb{D}}$, it follows that $W_{\mathbb{H}}(A) = \overline{\mathbb{B}}$. Let $\alpha \in W(\omega(A))$. Then, $\alpha = u^* \omega(A) u$, where $u \in \mathbb{C}^{2n}$ and $\|u\| = 1$. Decompose $u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$, where $u_1, u_2 \in \mathbb{C}^n$,

and define $w := \begin{bmatrix} -\bar{u}_2 \\ \bar{u}_1 \end{bmatrix}$. Note that

$$\begin{aligned} w^*u &= -\bar{u}_2^*u_1 + \bar{u}_1^*u_2 \\ &= -u_2^T u_1 + u_1^T u_2 \\ &= -(u_1^T u_2)^T + u_1^T u_2 = 0. \end{aligned}$$

Define $x := u_1 - ju_2 \in \mathbb{H}^n$. Then, $\|x\| = 1$ and for some $\beta \in \mathbb{C}$

$$\begin{aligned} \begin{bmatrix} \alpha & \beta \\ -\beta & \bar{\alpha} \end{bmatrix} &= \begin{bmatrix} u^* \\ w^* \end{bmatrix} \omega(A) \begin{bmatrix} u & w \end{bmatrix} \\ &= \begin{bmatrix} u_1^* & u_2^* \\ \bar{u}_2^* & \bar{u}_1^* \end{bmatrix} \omega(A) \begin{bmatrix} u_1 & -\bar{u}_2 \\ u_2 & \bar{u}_1 \end{bmatrix} = \omega(x)^* \omega(A) \omega(x). \end{aligned}$$

Hence, $\alpha + j\beta \in W_{\mathbb{H}}(A) = \overline{\mathbb{B}}$. Thus, $\|\alpha\|^2 + \|\beta\|^2 = \|\alpha + j\beta\|^2 \leq 1$, and therefore, $\|\alpha\| \leq 1$, which proves $\alpha \in \overline{\mathbb{D}}$. □

Before we can state and prove our main result, we require the following results. For the results that are known, we omit the proofs and provide only a reference.

Lemma 3.4 [2, Corollary 1] *Let $A \in \mathbb{H}^{n \times n}$. Then, $\mathbb{R} \cap W_{\mathbb{H}}(A)$ is a closed interval.*

Lemma 3.5 [2, Theorem 3] *For $A \in \mathbb{H}^{n \times n}$, $W_{\mathbb{H}}(A)$ is convex if and only if $\mathbb{R} \cap W_{\mathbb{H}}(A) = \{\text{Re}(q) : q \in W_{\mathbb{H}}(A)\}$.*

Corollary 3.6 *Let $A \in \mathbb{H}^{n \times n}$ and $W_{\mathbb{H}}(A) \subseteq \overline{\mathbb{B}}$. If $\pm 1 \in W_{\mathbb{H}}(A)$, then $W_{\mathbb{H}}(A)$ is convex.*

Proof Since $W_{\mathbb{H}}(A) \subseteq \overline{\mathbb{B}}$, we have

$$\{\text{Re}(q) : q \in W_{\mathbb{H}}(A)\} \subseteq [-1, 1].$$

As $\pm 1 \in \mathbb{R} \cap W_{\mathbb{H}}(A)$, it follows by Lemma 3.4 that:

$$[-1, 1] \subseteq \mathbb{R} \cap W_{\mathbb{H}}(A).$$

Then

$$[-1, 1] \subseteq \mathbb{R} \cap W_{\mathbb{H}}(A) \subseteq \{\text{Re}(q) : q \in W_{\mathbb{H}}(A)\} \subseteq [-1, 1].$$

Thus

$$\mathbb{R} \cap W_{\mathbb{H}}(A) = \{\text{Re}(q) : q \in W_{\mathbb{H}}(A)\},$$

and by Lemma 3.5 $W_{\mathbb{H}}(A)$ is convex. □

Our main result is the following description of quaternionic matrices whose bild is the unit disk.

Theorem 3.7 *Let $A \in \mathbb{H}^{n \times n}$. Then, $B(A) = \overline{\mathbb{D}}$ if and only if there exist $P_0, P_1 \in \mathbb{H}^{n-1 \times n}$, such that*

$$P_0^* P_0 + P_1^* P_1 = I_n \quad \text{and} \quad \frac{1}{2} A = P_0^* P_1.$$

Proof Assume there exist $P_0, P_1 \in \mathbb{H}^{n-1 \times n}$, such that

$$P_0^* P_0 + P_1^* P_1 = I_n \quad \text{and} \quad \frac{1}{2} A = P_0^* P_1.$$

Then

$$\omega(P_0)^* \omega(P_0) + \omega(P_1)^* \omega(P_1) = I_{2n} \quad \text{and} \quad \frac{1}{2} \omega(A) = \omega(P_0)^* \omega(P_1).$$

Since $\omega(P_0), \omega(P_1) \in \mathbb{C}^{2(n-1) \times 2n}$, it follows by [16, Theorem 4.5] that $W(\omega(A)) = \overline{\mathbb{D}}$, and hence, by Lemma 3.1, we have $B(A) \subseteq \overline{\mathbb{D}}$. From [2, Theorem 1], it follows:

$$W_{\mathbb{H}}(A) \subseteq \{a_0 + p : a \in \mathbb{R}, p \in \mathbb{H}, \operatorname{Re}(p) = 0, a_0 + \|p\|i \in B(A)\}.$$

Therefore, it is clear that $W_{\mathbb{H}}(A) \subseteq \overline{\mathbb{B}}$.

Next, note that

$$I_n - \frac{1}{2} A - \frac{1}{2} A^* = (P_0 - P_1)^* (P_0 - P_1) \geq 0,$$

where $P_0 - P_1 \in \mathbb{H}^{n-1 \times n}$. Hence, $\operatorname{Ker}(P_0 - P_1)$ is non-trivial. Take $x \in \mathbb{H}^n$ where $x \in \operatorname{Ker}(P_0 - P_1)$ and $\|x\| = 1$. Then

$$\begin{aligned} 0 &= x^* (P_0 - P_1)^* (P_0 - P_1) x \\ &= x^* \left(I_n - \frac{1}{2} A - \frac{1}{2} A^* \right) x \\ &= x^* x - \frac{1}{2} x^* A x - \frac{1}{2} x^* A^* x \\ &= 1 - \operatorname{Re}(x^* A x). \end{aligned}$$

Since $x^* A x \in W_{\mathbb{H}}(A)$, it follows that $\|x^* A x\| \leq 1$, and hence, $1 = x^* A x \in W_{\mathbb{H}}(A)$. In a similar way, it can be proven that $-1 \in W_{\mathbb{H}}(A)$ using

$$I_n + \frac{1}{2} A + \frac{1}{2} A^* = (P_0 + P_1)^* (P_0 + P_1) \geq 0.$$

Hence, by Corollary 3.6, it follows that $W_{\mathbb{H}}(A)$ is convex, and therefore, $B(A) = W(\omega(A)) = \overline{\mathbb{D}}$ by Lemma 3.3.

For the other direction, assume $B(A) = \overline{\mathbb{D}}$. By Lemma 3.3, it then follows that $B(A) = W(\omega(A))$. Thus, $|x^* \omega(A)x| \leq 1$ for all $x \in \mathbb{C}^{2n}$ where $\|x\| = 1$. This is equivalent to

$$\operatorname{Re} \left(e^{i\theta} x^* \omega(A)x \right) \leq 1, \quad \text{for all } \theta \text{ and } \|x\| = 1,$$

which can be rewritten as

$$\frac{1}{2} x^* \left(e^{i\theta} \omega(A) + e^{-i\theta} \omega(A)^* \right) x \leq x^* x, \quad \text{for all } \theta \text{ and } \|x\| = 1.$$

Hence

$$I_{2n} - \frac{1}{2} \left(e^{i\theta} \omega(A) + e^{-i\theta} \omega(A)^* \right) \geq 0 \quad \text{for all } \theta,$$

or equivalently

$$I_{2n} - \frac{1}{2} \left(z \omega(A) + \frac{1}{z} \omega(A)^* \right) \geq 0 \quad \text{for all } z \in \mathbb{T}.$$

Define $Q_0 := I_{2n}$ and $Q_1 := -\frac{1}{2} \omega(A)$. Then, $Q_j \in \Omega_{2n}$ for $j = -1, 0, 1$, where $Q_{-1} := Q_1^*$, and

$$Q(z) := \sum_{j=-1}^1 Q_j z^j \geq 0 \quad \text{for all } z \in \mathbb{T}.$$

By the Fejér–Riesz factorization (Theorem 2.3), there exist $\tilde{P}_0, \tilde{P}_1 \in \Omega_{2n}$, such that

$$Q(z) = (\tilde{P}_0 + z \tilde{P}_1)^* (\tilde{P}_0 + z \tilde{P}_1), \quad \text{for all } z \in \mathbb{T},$$

and in fact, we choose the factorization that corresponds to the choice X_{\min} in Theorem 2.3. By equating coefficients, it follows that:

$$P_0^* P_0 + P_1^* P_1 = I_n \quad \text{and} \quad \frac{1}{2} A = P_0^* P_1,$$

where $\omega(P_0) = \tilde{P}_0$ and $\omega(P_1) = \tilde{P}_1$, for some $P_0, P_1 \in \mathbb{H}^{n \times n}$. Hence, it remains to prove that we are able to reduce the size of P_0 and P_1 .

Note that since $\bar{z} \in W(\omega(A))$ for any $z \in \mathbb{T}$, there exists a $y \in \mathbb{C}^{2n}$, $\|y\| = 1$, such that $\bar{z} = y^* \omega(A) y$. Then

$$\begin{aligned} y^* (\tilde{P}_0 + z\tilde{P}_1)^* (\tilde{P}_0 + z\tilde{P}_1) y &= y^* Q(z) y \\ &= y^* \left(I_{2n} - \frac{1}{2} z \omega(A) - \frac{1}{2} \frac{1}{z} \omega(A)^* \right) y = 0, \end{aligned}$$

which proves that $\tilde{P}_0 + z\tilde{P}_1$ is singular for all $z \in \mathbb{T}$. We reason by contradiction that \tilde{P}_0 is singular. If \tilde{P}_0 is non-singular, then

$$\tilde{P}_0 + z\tilde{P}_1 = \tilde{P}_0 \left(I_{2n} + z\tilde{P}_0^{-1}\tilde{P}_1 \right) = \tilde{P}_0 \left(I_{2n} - z \left(-\tilde{P}_0^{-1}\tilde{P}_1 \right) \right).$$

Using that

$$\det \left(I_{2n} - z \left(-\tilde{P}_0^{-1}\tilde{P}_1 \right) \right) = 0$$

only holds when $\frac{1}{z}$ is an eigenvalue of $-\tilde{P}_0^{-1}\tilde{P}_1$, of which there are at most $2m$, we arrive at a contradiction. Hence, \tilde{P}_0 is singular. By a similar argument, it follows that \tilde{P}_1 is singular.

Write

$$0 \leq \begin{bmatrix} \tilde{P}_0 \\ \tilde{P}_1 \end{bmatrix}^* \begin{bmatrix} \tilde{P}_0 & \tilde{P}_1 \end{bmatrix} =: \begin{bmatrix} P & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P \end{bmatrix},$$

where we recall that we made the choice corresponding to X_{\min} in Theorem 2.3. This results in choosing the largest P in the Loewner order, which we denote by P_{\max} . Furthermore

$$\text{rank} \begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} \end{bmatrix} = \text{rank } P_{\max}.$$

To see this, we use the generalized Schur complement of P_{\max} which equals

$$B := I_{2n} - P_{\max} - \frac{1}{2}\omega(A)^* P_{\max}^{(-1)} \frac{1}{2}\omega(A) \geq 0,$$

with $P_{\max}^{(-1)}$ the Moore–Penrose generalized inverse of P_{\max} . From

$$\text{rank} \begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} \end{bmatrix} = \text{rank } P_{\max} + \text{rank } B,$$

it follows that

$$\text{rank} \begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} \end{bmatrix} \geq \text{rank } P_{\max}.$$

Suppose

$$\text{rank} \begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} \end{bmatrix} > \text{rank } P_{\max}.$$

Then, $B \neq 0$. Furthermore,

$$\begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} - B \end{bmatrix} = \begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & \frac{1}{2}\omega(A)^* P_{\max}^{(-1)} \frac{1}{2}\omega(A) \end{bmatrix} \geq 0.$$

Now

$$\begin{bmatrix} P_{\max} + B & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - (P_{\max} + B) \end{bmatrix} \geq 0,$$

and since $P_{\max} + B > P_{\max}$, this is a contradiction.

Let

$$k = \text{rank } P_{\max} = \text{rank} \begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} \end{bmatrix}.$$

Since $P_{\max} = \tilde{P}_0^* \tilde{P}_0$ and $\text{rank } \tilde{P}_0 < 2n$, we have that $k < 2n$. Since 0 as an eigenvalue of P_{\max} has even multiplicity (see, e.g., [21, Corollary 5.1]), we have that $k \leq 2(n-1)$. Then, by the spectral theorem in quaternions [9, Proposition 3.7], there exist $R_0, R_1 \in \Omega_{2(n-1) \times 2n}$, such that

$$\begin{bmatrix} P_{\max} & \frac{1}{2}\omega(A) \\ \frac{1}{2}\omega(A)^* & I_{2n} - P_{\max} \end{bmatrix} = \begin{bmatrix} R_0 \\ R_1 \end{bmatrix}^* \begin{bmatrix} R_0 & R_1 \end{bmatrix}.$$

Now, $S_0 = \omega^{-1}(R_0) \in \mathbb{H}^{(n-1) \times n}$, $S_1 = \omega^{-1}(R_1) \in \mathbb{H}^{(n-1) \times n}$ gives us the desired matrices, so that $A = 2S_0^* S_1$ and $S_0^* S_0 + S_1^* S_1 = I_n$. □

Let us illustrate Theorem 3.7 in a simple example.

Example 3.8 Let $A = \begin{bmatrix} 0 & a_1 & 0 \\ 0 & 0 & a_2 \\ 0 & 0 & 0 \end{bmatrix}$. Then, one easily checks that $B(A)$ is a disk with center 0 (see also [5]). In this case, let

$$P_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{a_2}{2} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad P_1 = \begin{pmatrix} 0 & \frac{a_1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Then, $A = 2P_0^* P_1$. Moreover, when $\|a_1\|^2 + \|a_2\|^2 = 4$ we obtain that $P_0^* P_0 + P_1^* P_1 = I_3$. Therefore, under this condition, we have that the numerical range of A is the closed unit ball in \mathbb{H} .

Acknowledgements The authors gratefully acknowledge funding from the National Graduate Academy for Mathematical and Statistical Sciences (NGA(MaSS)). The research of HJW is supported by National Science Foundation under Grant DMS 2000037.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Aghamollaei, G., Aboutalebi, N.H., Kermani, H.M.: Some results on the k -numerical range of quaternion matrices. *Linear Multilinear Algebra* **64**(12), 2419–2430 (2016)
2. Au-Yeung, Y.H.: On the convexity of numerical range in quaternionic Hilbert spaces. *Linear Multilinear Algebra* **16**(1–4), 93–100 (1984)
3. Au-Yeung, Y.H.: A short proof of a theorem on the numerical range of a normal quaternionic matrix. *Linear Multilinear Algebra* **39**(3), 279–284 (1995)
4. Bakonyi, M., Woerdeman, H.J.: *Matrix Completions, Moments, and Sums of Hermitian Squares*. Princeton University Press, Princeton (2011)
5. Carvalho, L., Diogo, C., Mendes, S.: On the convexity and circularity of the numerical range of nilpotent quaternionic matrices. *New York J. Math.* **25**, 1385–1404 (2019)
6. Chien, M.T., Nakazato, H.: Boundary generating curves of quaternionic numerical ranges. *Linear Algebra Appl.* **574**, 201–215 (2019)
7. Daubechies, I.: *Ten Lectures on Wavelets* (CBMS-NSF Regional Conference Series in Applied Mathematics). SIAM: Society for Industrial and Applied Mathematics, Philadelphia (1992)
8. Dritschel, M., Woerdeman, H.J.: Outer factorizations in one and several variables. *Trans. Am. Math. Soc.* **357**, 4661–4679 (2005)
9. Farenick, D.R., Pidkowich, B.A.F.: The spectral theorem in quaternions. *Linear Algebra Appl.* **371**, 75–102 (2003)
10. Francis, B.A.: *A Course in H^∞ Control Theory*. Lecture Notes in Control and Information Sciences, vol. 88, Corrected edn. Springer, Berlin (1987)
11. Geronimo, J.S., Woerdeman, H.J.: Positive extensions, Fejér–Riesz factorization and autoregressive filters in two variables. *Ann. Math.* **160**, 839–906 (2004)
12. Hachez, Y., Woerdeman, H.J.: The Fischer–Frobenius transformation and outer factorization. In: *Operator Theory, Structured Matrices, and Dilations*. Editors: Mihaly Bakonyi, Aurelian Gheondea, Mihai Putinar, and James Rovnyak. Theta Series in Advanced Mathematics, vol. 7, pp. 181–203. Theta, Bucharest (2007)
13. Hausdorff, F.: Der Wertvorrat einer Bilinearform. (German) *Math. Z.* **3**(1), 314–316 (1919)
14. Kippenhahn, R.: Über den Wertvorrat einer Matrix. (German) *Math. Nachr.* **6**, 193–228 (1951). [English translation: On the numerical range of a matrix. Translated from German by Paul F. Zachlin and Michiel E. Hochstenbach. *Linear Multilinear Algebra* **56**, 185–225 (2008)]
15. Lancaster, P., Rodman, L.: *Algebraic Riccati Equations*. Oxford Science Publications, The Clarendon Press, Oxford University Press, New York (1995)
16. Poon, E., Spitkovsky, I.M., Woerdeman, H.J.: Factorization of singular matrix polynomials and matrices with circular higher rank numerical ranges. *SIAM J. Matrix Anal. Appl.* **43**(3), 1423–1439 (2022)
17. Ramesh, G.: On the numerical radius of a quaternionic normal operator. *Adv. Oper. Theory* **2**(1), 78–86 (2017)
18. Rodman, L.: *Topics in Quaternion Linear Algebra*. Princeton Series in Applied Mathematics. Princeton University Press, Princeton (2014)
19. So, W., Thompson, R.C.: Convexity of the upper complex plane part of the numerical range of a quaternionic matrix. *Linear Multilinear Algebra* **41**, 303–365 (1996)

-
20. Toeplitz, O.: Das algebraische Analogon zu einem Satze von Fejér. (German) *Math. Z.* **2**(1–2), 187–197 (1918)
 21. Zhang, F.: Quaternions and matrices of quaternions. *Linear Algebra Appl.* **251**, 21–57 (1997)